

B4

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 May 2002 (10.05.2002)

PCT

(10) International Publication Number
WO 02/36584 A1

(51) International Patent Classification⁷: C07D 333/54, 275/04, 409/12, 417/12, A61K 31/381, A61P 3/00

(74) Agent: WITTE, Hubert; 124 Grenzacherstrasse, CH-4070 Basle (CH).

(21) International Application Number: PCT/EP01/12451

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: 26 October 2001 (26.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00123826.0 2 November 2000 (02.11.2000) EP

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; 124 Grenzacherstrasse, CH-4070 Basle (CH).

(72) Inventors: AEBI, Johannes; 13, Wallstrasse, CH-4051 Basle (CH). ACKERMANN, Jean; 16, Im Glögglihof, CH-4125 Riehen (CH). DEHMLow, Henrietta; 85, Ritterstrasse, 79639 Grenzach-Wyhlen (DE). MAERKI, Hans-Peter; Seltisbergerstrasse 75, CH-4059 Basle (CH). MORAND, Olivier; 2, rue de Vignes, F-68220 Hegenheim (FR).


Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/36584 A1

(54) Title: CHOLESTEROL LOWERING BENZO'B' THIOPHENES AND BENZO'D' ISOTHIAZOLES




(57) Abstract: The present invention relates to compounds of formula (I) wherein U, A¹, A², A³, A⁴, V, X, Y, Z, Q, m and n are as defined in the description and claims and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof. The compounds are useful for the treatment and/or prophylaxis of diseases which are associated with 2,3-oxidosqualene-lanosterol cyclase such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, gallstones, tumors and/or hyperproliferative

disorders, and treatment and/or prophylaxis of impaired glucose tolerance and diabetes.

CHOLESTEROL LOWERING BENZO[B]THIOPHENES AND BENZO[D]ISOTHIAZOLES

The present invention is concerned with novel heteroaromates, their manufacture and their use as medicaments. In particular, the invention relates to compounds of the formula (I)



5 wherein

U is O or a lone pair,

Y N, or CR¹,

Z is S, or S(O₂),

V is a) -CH=CH-, and m and n are 0, or

10 b) -CH₂-, and m+n ≤ 2, or

c) O or NR², and m is 1 to 6, n is 1 to 6, m+n ≤ 7, or m is 1 to 3, n is 0, or

d) S, and m is 1 to 7, n is 0 to 6, m+n ≤ 7, or

e) -C≡C-, and m is 0 to 7, n is 0 to 7, m+n is ≤ 7,

15 Q is cycloalkyl, cycloalkyl-lower-alkyl, phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R³, or an alkyl, alkenyl or alkadienyl group optionally substituted by OH,

A¹ is hydrogen, lower-alkenyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy,

20 A² is cycloalkyl, cycloalkyl-lower-alkyl, lower-alkenyl, lower-alkinyl, heteroaryl, or lower-alkyl optionally substituted with halogen, hydroxy, lower-alkoxy, or thio-lower-alkoxy, or

A¹ and A² are bonded to each other to form a ring and -A¹-A²- is lower-alkylene, or lower-alkenylene, optionally substituted by R⁴, in which one -CH₂- group of

- 2 -

- A¹-A²- can optionally be replaced by a NR⁵, S, or O, or -A¹-A²- is
 -CH=N=CH=CH- which can optionally be substituted by lower-alkyl,
 A³ and A⁴ independently from each other are hydrogen or lower-alkyl, or
 A³ and A⁴ are bonded to each other to form a ring together with the carbon atom to which
 5 they are attached and -A³-A⁴- is -(CH₂)₂₋₅- which can optionally be mono- or
 multiply-substituted by lower-alkyl,
 X is hydrogen or one or more optional halogen and/or lower-alkyl substituents,
 R¹ is hydrogen or lower-alkyl,
 R² is hydrogen or lower-alkyl,
 10 R³ is halogen, N(R⁶,R⁷), piperidyl, piperazinyl, piperazinyl-ethanone,
 morpholinyl, CONH₂, CN, NO₂, CF₃, OH, lower-alkoxy, thio-lower-
 alkoxy, or is lower-alkyl, lower-alkenyl, or lower-alkinyl, optionally
 substituted with OH, SH or N(R⁸,R⁹),
 R⁴ is hydroxy, lower-alkyl, lower-alkoxy, or thio-lower-alkoxy,
 15 R⁵ is hydrogen, lower-alkyl, or lower-alkyl-carbonyl,
 R⁶ and R⁷ independently from each other are selected from the group consisting of
 hydrogen, lower-alkyl, phenyl and benzyl,
 R⁸ and R⁹ independently from each other are selected from the group consisting of
 hydrogen and lower-alkyl,
 20 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

The compounds of the present invention inhibit 2,3-oxidosqualene-lanosterol
 cyclase (EC 5.4.99.) which is required for the biosynthesis of cholesterol, ergosterol and
 other sterols. Causal risk factors that directly promote the development of coronary and
 peripheral atherosclerosis include elevated low-density lipoprotein cholesterol (LDL-C),
 25 low high-density lipoprotein cholesterol (HDL-C), hypertension, cigarette smoking and
 diabetes mellitus. Other synergistic risk factors include elevated concentrations of
 triglyceride (TG)-rich lipoproteins, small, dense low-density lipoprotein particles,
 lipoprotein (a) (Lp(a)), and homocysteine. Predisposing risk factors modify the causal or
 conditional risk factors and thus affect atherogenesis indirectly. The predisposing risk
 30 factors are obesity, physical inactivity, family history of premature CVD, and male sex. The
 strong connection between coronary heart disease (CHD) and high LDL-C levels in
 plasma, and the therapeutic advantage of lowering elevated LDL-C levels are now well
 established (Gotto et al., Circulation 81, 1990, 1721-1733; Stein et al., Nutr. Metab.

Cardiovasc. Dis. 2, 1992, 113-156; Illingworth, Med. Clin. North. Am. 84, 2000, 23-42). Cholesterol-rich, sometimes unstable, atherosclerotic plaques lead to the occlusion of blood vessels resulting in an ischemia or an infarct. Studies with respect to primary prophylaxis have shown that a lowering of plasma LDL-C levels in plasma reduces the frequency of non-fatal incidences of CHD, while the overall morbidity remains unchanged. The lowering of plasma LDL-C levels in patients with pre-established CHD (secondary intervention) reduces CHD mortality and morbidity; meta-analysis of different studies shows that this decrease is proportional to the reduction of the LDL-C (Ross et al., Arch. Intern. Med. 159, 1999, 1793-1802).

10 The clinical advantage of cholesterol lowering is greater for patients with pre-established CHD than for asymptomatic persons with hypercholesterolemia. According to current guidelines, cholesterol lowering treatment is recommended for patients who had survived a myocardial infarct or patients suffering from angina pectoris or another atherosclerotic disease, with a target LDL-C level of 100 mg/dl.

15 Preparations such as bile acid sequestrants, fibrates, nicotinic acid, probucol as well as statins, i.e. HMG-Co-A reductase inhibitors such as simvastatin and atorvastatin, are used for usual standard therapies. The best statins reduce plasma LDL-C effectively by at least 40%, and also plasma triglycerides, a synergistic risk factor, but less effectively. In contrast, fibrates reduce plasma triglycerides effectively, but not LDL-C. Combination of a statin and a fibrate proved to be very efficacious in lowering LDL-C and triglycerides (Ellen and McPherson, J. Cardiol. 81, 1998, 60B-65B), but safety of such a combination remains an issue (Shepherd, Eur. Heart J. 16, 1995, 5-13). A single drug with a mixed profile combining effective lowering of both LDL-C and triglycerides would provide additional clinical benefit to asymptomatic and symptomatic patients.

25 In humans, statins are well tolerated at standard dosage, but reductions in non-sterol intermediates in the cholesterol synthesis pathway, such as isoprenoids and coenzyme Q, may be associated with adverse clinical events at high doses (Davignon et al., Can. J. Cardiol. 8, 1992, 843-864; Pederson and Tobert, Drug Safety 14, 1996, 11-24).

This has stimulated the search for, and development of compounds that inhibit cholesterol biosynthesis, yet act distal to the synthesis of these important, non-sterol intermediates. 2,3-oxidosqualene:lanosterol cyclase (OSC), a microsomal enzyme, represents a unique target for a cholesterol-lowering drug (Morand et al., J. Lipid Res., 38, 1997, 373-390; Mark et al., J. Lipid Res. 37, 1996, 148-158). OSC is downstream of farnesyl-pyrophosphate, beyond the synthesis of isoprenoids and coenzyme Q. In hamsters, pharmacologically active doses of an OSC inhibitor showed no adverse side-effects, in contrast to a statin which reduced food-intake and body weight, and increased

plasma bilirubin, liver weight and liver triglyceride content (Morand et al., J. Lipid Res., 38, 1997, 373-390). The compounds described in European Patent Application No. 636 367, which inhibit OSC and which lower the total cholesterol in plasma, belong to these substances.

5 OSC inhibition does not trigger the overexpression of HMGR because of an indirect, negative feed-back regulatory mechanism involving the production of 24(S),25-epoxycholesterol (Peffley et al., Biochem. Pharmacol. 56, 1998, 439-449; Nelson et al., J. Biol. Chem. 256, 1981, 1067-1068; Spencer et al., J. Biol. Chem. 260, 1985, 13391-13394; Panini et al., J. Lipid Res. 27, 1986, 1190-1204; Ness et al., Arch. Biochem. Biophys. 308, 10 1994, 420-425). This negative feed-back regulatory mechanism is fundamental to the concept of OSC inhibition because (i) it potentiates synergistically the primary inhibitory effect with an indirect down-regulation of HMGR, and (ii) it prevents the massive accumulation of the precursor monooxidosqualene in the liver. In addition, 24(S),25-epoxycholesterol was found to be one of the most potent agonists of the nuclear receptor 15 LXR (Janowski et al., Proc. Natl. Acad. Sci. USA, 96, 1999, 266-271). Considering that 24(S),25-epoxycholesterol is a by-product of inhibition of OSC it is hypothesized that the OSC inhibitors of the present invention could also indirectly activate LXR-dependent pathways such as (i) cholesterol-7 α -hydroxylase to increase the consumption of cholesterol via the bile acid route, (ii) expression of ABC proteins with the potential to 20 stimulate reverse cholesterol transport and increase plasma HDL-C levels (Venkateswaran et al., J. Biol. Chem. 275, 2000, 14700-14707; Costet et al., J. Biol. Chem. June 2000, in press; Ordovas, Nutr Rev 58, 2000, 76-79, Schmitz and Kaminsky, Front Biosci 6, 2001, D505-D514), and/or inhibit intestinal cholesterol absorption (Mangelsdorf, XIIth International Symposium on Atherosclerosis, Stockholm, June 2000). In addition, possible 25 cross talks between fatty acid and cholesterol metabolism mediated by liver LXR have been hypothesized (Tobin et al., Mol. Endocrinol. 14, 2000, 741-752).

The present compounds of formula I inhibit OSC and therefore also inhibit the biosynthesis of cholesterol, ergosterol and other sterols, and reduce the plasma cholesterol levels. They can therefore be used in the therapy and prophylaxis of hypercholesterolemia, 30 hyperlipemia, arteriosclerosis and vascular diseases in general. Furthermore, they can be used in the therapy and/or prevention of mycoses, parasite infections, gallstones, cholestatic liver disorders, tumors and hyperproliferative disorders, e.g. hyperproliferative skin and vascular disorders. In addition, it has unexpectedly been found that the compounds of the present invention can also be of therapeutical use to improve glucose 35 tolerance in order to treat and/or prevent related diseases such as diabetes. The compounds of the present invention further exhibit improved pharmacological properties compared to known compounds.

Unless otherwise indicated the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

In this specification the term "lower" is used to mean a group consisting of one to seven, preferably of one to four carbon atom(s).

5 The term "lone pair" refers to an unbound electron pair, in particular to the unbound electron pair of a nitrogen atom in e.g. an amine.

The term "halogen" refers to fluorine, chlorine, bromine and iodine, with fluorine, chlorine and bromine being preferred.

10 The term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon atoms. Lower-alkyl groups as defined below also are preferred alkyl groups. Alkyl groups can be substituted e.g. with halogen, particularly with fluorine or chlorine, hydroxy, lower-alkoxy, and/or lower-alkoxy-carbonyl.

15 The term "lower-alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent alkyl radical of one to seven carbon atoms, preferably one to four carbon atoms. This term is further exemplified by such radicals as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, t-butyl and the like. A lower-alkyl group may have a substitution pattern as described earlier in connection with the term
20 "alkyl".

The term "cycloalkyl" refers to a monovalent carbocyclic radical of 3 to 10 carbon atom(s), preferably 3 to 6 carbon atoms.

The term "alkoxy" refers to the group $R'-O-$, wherein R' is an alkyl. The term "lower-alkoxy" refers to the group $R'-O-$, wherein R' is a lower-alkyl. The term "thio-alkoxy"
25 refers to the group $R'-S-$, wherein R' is an alkyl. The term "thio-lower-alkoxy" refers to the group $R'-S-$, wherein R' is a lower-alkyl.

The term "alkenyl", alone or in combination with other groups, stands for a straight-chain or branched hydrocarbon residue comprising an olefinic bond and up to 20, preferably up to 16 carbon atoms, more preferably up to 10 carbon atoms. Lower-alkenyl
30 groups as described below also are preferred alkenyl groups. The term "lower-alkenyl" refers to a straight-chain or branched hydrocarbon residue comprising an olefinic bond and up to 7, preferably up to 4 carbon atoms, such as e.g. 2-propenyl. An alkenyl or lower-alkenyl group may have a substitution pattern as described earlier in connection with the

term "alkyl".

The term "alkadienyl", alone or in combination with other groups, stands for a straight-chain or branched hydrocarbon residue comprising 2 olefinic bonds and up to 20, preferably up to 16 carbon atoms, more preferably up to 10 carbon atoms. Lower-
5 alkadienyl groups as described below also are preferred alkadienyl groups. The term "lower-alkadienyl" refers to a straight-chain or branched hydrocarbon residue comprising 2 olefinic bonds and up to 7 carbon atoms. An alkadienyl or lower-alkadienyl group may have a substitution pattern as described earlier in connection with the term "alkyl".

The term "alkinyl", alone or in combination with other groups, stands for a straight-
10 chain or branched hydrocarbon residue comprising a tripple bond and up to 20, preferably up to 16 carbon atoms. The term "lower-alkinyl" refers to a straight-chain or branched hydrocarbon residue comprising a tripple bond and up to 7, preferably up to 4 carbon atoms, such as e.g. 2-propinyl. An alkinyl or lower-alkinyl group may have a substitution pattern as described earlier in connection with the term "alkyl".

15 The term "alkylene" refers to a straight chain or branched divalent saturated aliphatic hydrocarbon group of 1 to 20 carbon atoms, preferably 1 to 16 carbon atoms. The term "lower-alkylene" refers to a straight chain or branched divalent saturated aliphatic hydrocarbon group of 1 to 7, preferably 3 to 6 carbon atoms. An alkylene or lower-alkylene group may have a substitution pattern as described earlier in connection with the
20 term "alkyl".

The term "alkenylene" refers to a straight chain or branched divalent hydrocarbon group comprising an olefinic bond and up to 20 carbon atoms, preferably up to 16 carbon atoms. The term "lower-alkenylene" refers to a straight chain or branched divalent hydrocarbon group comprising an olefinic bond and up to 7, preferably up to 6 C-atoms.
25 An alkenylene or lower-alkenylene group may have a substitution pattern as described earlier in connection with the term "alkyl".

The term "aryl" relates to the phenyl or naphthyl group which can optionally be mono- or multiply-substituted by lower-alkyl, dioxo-lower-alkylene (forming e.g. a benzodioxyl group), halogen, hydroxy, cyano, CF₃, NH₂, N(lower-alkyl)₂, aminocarbonyl,
30 carboxy, nitro, lower-alkoxy, thio-lower-alkoxy, lower-alkylcarbonyl, lower-alkylcarbonyloxy, aryl, or aryloxy. Preferred substituents are lower-alkyl, lower-alkoxy, thio-lower-alkoxy, lower-alkyl-carbonyl, lower-alkoxycarbonyl, fluorine, chlorine, bromine, CN, CF₃, and/or dioxo-lower-alkylene. More preferred substituents are fluorine, chlorine, bromine and CF₃.


35 The term "heteroaryl" refers to an aromatic 5- or 6-membered ring which can

comprise 1, 2 or 3 atoms selected from nitrogen, oxygen and/or sulphur such as furyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, thienyl, isoxazolyl, oxazolyl, imidazolyl, or pyrrolyl. A heteroaryl group may have a substitution pattern as described earlier in connection with the term "aryl".

- 5 The term "pharmaceutically acceptable salts" embraces salts of the compounds of formula (I) with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulphuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic acid, fumaric acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like, which are non toxic to living organisms. Preferred salts are formates,
10 hydrochlorides and hydrobromides.

- The term "pharmaceutically acceptable esters" embraces esters of the compounds of formula (I), in which hydroxy groups have been converted to the corresponding esters with inorganic or organic acids such as nitric acid, sulphuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic acid, succinic acid, tartaric acid, methanesulphonic
15 acid, p-toluenesulphonic acid and the like, which are non toxic to living organisms.

In detail, the present invention relates to compounds of formula (I)



wherein

- U is O or a lone pair,
- 5 Y N, or CR¹,
- Z is S, or S(O₂),
- V is a) -CH=CH-, and m and n are 0, or
 b) -CH₂-, and m+n ≤ 2, or
 c) O or NR², and m is 1 to 6, n is 1 to 6, m+n ≤ 7, or m is 1 to 3, n is 0, or
 10 d) S, and m is 1 to 7, n is 0 to 6, m+n ≤ 7, or
 e) -C≡C-, and m is 0 to 7, n is 0 to 7, m+n is ≤ 7,
- Q is cycloalkyl, cycloalkyl-lower-alkyl, phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R³, or an alkyl, alkenyl or alkadienyl group optionally substituted by OH,
- 15 A¹ is hydrogen, lower-alkenyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy,
- A² is cycloalkyl, cycloalkyl-lower-alkyl, lower-alkenyl, lower-alkinyl, heteroaryl, or lower-alkyl optionally substituted with halogen, hydroxy, lower-alkoxy, or thio-lower-alkoxy, or
- 20 A¹ and A² are bonded to each other to form a ring and -A¹-A²- is lower-alkylene, or lower-alkenylene, optionally substituted by R⁴, in which one -CH₂- group of -A¹-A²- can optionally be replaced by a NR⁵, S, or O, or -A¹-A²- is -CH=N=CH=CH- which can optionally be substituted by lower-alkyl,
- A³ and A⁴ independently from each other are hydrogen or lower-alkyl, or
- 25 A³ and A⁴ are bonded to each other to form a ring together with the carbon atom to which they are attached and -A³-A⁴- is -(CH₂)₂₋₅- which can optionally be mono- or multiply-substituted by lower-alkyl,

- X is hydrogen or one or more optional halogen and/or lower-alkyl substituents,
- R¹ is hydrogen or lower-alkyl,
- R² is hydrogen or lower-alkyl,
- R³ is halogen, N(R⁶, R⁷), piperidyl, piperazinyl, piperazinyl-ethanone,
 5 morpholinyl, CONH₂, CN, NO₂, CF₃, OH, lower-alkoxy, thio-lower-alkoxy, or is lower-alkyl, lower-alkenyl, or lower-alkinyl, optionally substituted with OH, SH or N(R⁸, R⁹),
- R⁴ is hydroxy, lower-alkyl, lower-alkoxy, or thio-lower-alkoxy,
- R⁵ is hydrogen, lower-alkyl, or lower-alkyl-carbonyl,
- 10 R⁶ and R⁷ independently from each other are selected from the group consisting of hydrogen, lower-alkyl, phenyl and benzyl,
- R⁸ and R⁹ independently from each other are selected from the group consisting of hydrogen and lower-alkyl,

and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

- 15 Preferred are compounds of formula (I) and/or pharmaceutically acceptable salts thereof. Other preferred embodiments relate to compounds of formula (I) wherein U is a lone pair or to compounds of formula (I) wherein U is O.

Each of the definitions of V given above, a), b), c), d) and e), individually or in combination constitutes a preferred embodiment of the present invention. Compounds as
 20 described above in which V is O, m is 1 to 6, n is 1 to 6, and m+n ≤ 7 relate to a further preferred embodiment of the present invention. Another preferred embodiment of the present invention are compounds as described above, wherein V is O, m is 1 to 3, and n is 0. Other preferred compounds are those, wherein V is -C≡C-, m is 0 to 2, and n is 0. Further preferred compounds are those, wherein V is -CH₂- and m+n ≤ 2.

- 25 Other preferred compounds of the present invention are those in which A¹ represents hydrogen, methyl, or ethyl optionally substituted with hydroxy or methoxy. Another group of preferred compounds of the present invention are those in which A² represents methyl, n-propyl, i-propyl, 2-propenyl, 2-propinyl, 1,1-dimethyl-2-propinyl, cyclopropyl-methylene, or ethyl optionally substituted with hydroxy or methoxy, with
 30 those compounds wherein A² represents methyl, ethyl, 2-hydroxy-ethyl, 2-methoxy-ethyl, n-propyl, 2-propenyl, or cyclopropyl-methylene being especially preferred. If A² is heteroaryl, pyridyl and pyrimidinyl, which can optionally be substituted with lower-alkyl, preferably with methyl, are preferred heteroaryl groups. Further preferred compounds as

defined above are those wherein A^2 is 2-Methyl-pyrimidin-4-yl.

Compounds of formula (I), wherein A^1 and A^2 are bonded to each other to form a ring and $-A^1-A^2-$ is lower-alkylene, optionally substituted by R^4 , in which one $-CH_2-$ group of $-A^1-A^2-$ can optionally be replaced by NR^5 or O, wherein R^4 is hydroxy, and R^5 is lower-alkyl or lower-alkyl-carbonyl, or $-A^1-A^2-$ is $-CH=N=CH=CH-$, are also preferred, with those compounds wherein said optional substituent R^5 is methyl being particularly preferred. If $-A^1-A^2-$ is $-CH=N=CH=CH-$ which can optionally be substituted by lower-alkyl, $-CH=N=CH=CH-$ is preferably substituted by methyl. In compounds wherein $-A^1-A^2-$ is $-CH=N=CH=CH-$, A^1 and A^2 , together with the nitrogen atom to which they are bound, form an imidazol ring. In compounds wherein A^1 and A^2 are bonded to each other to form a ring, said ring is preferably a 4-, 5-, or 6-membered ring such as e.g. piperidinyl or pyrrolidinyl. Compounds, wherein $-A^1-A^2-$ is $-(CH_2)_5-$ are also preferred.

A further preferred embodiment of the present invention relates to compounds of formula (I), wherein A^3 and/or A^4 represent hydrogen. In addition, compounds of formula (I) as defined above, wherein A^3 and/or A^4 represent methyl, relate to a preferred embodiment of the present invention. Compounds of formula (I) as defined above, wherein A^3 and A^4 are bonded to each other to form a cyclopentyl-ring or a cyclohexyl-ring together with the carbon atom to which they are attached and $-A^3-A^4-$ is $-(CH_2)_4-$ or $-(CH_2)_5-$ respectively represent another preferred embodiment of the present invention. The term $-(CH_2)_{2-5}-$ denotes the groups $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$ and $-(CH_2)_5-$.


In addition, compounds as described before, wherein Y is CR^1 and R^1 is methyl, are preferred. Furthermore, compounds in which Y is N relate to a preferred embodiment of the present invention. Other preferred compounds are those, wherein Z is S. Compounds, wherein Z is SO_2 are also preferred.

Compounds of formula (I), wherein Q is phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R^3 , or an alkyl or alkenyl group optionally substituted by OH, wherein R^3 is as defined above, represent a preferred embodiment of the present invention, with those compounds wherein Q is phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R^3 , wherein R^3 is fluorine, chlorine, bromine, or CF_3 being more preferred, and with those compounds wherein Q is 4-chloro-phenyl, 4-bromo-phenyl, or 4-trifluoromethyl-phenyl being particularly preferred. Another preferred group relates to compounds wherein X is hydrogen.

Further preferred embodiments of the present invention are those compounds as defined above wherein V is not $-CH_2-$ and/or not $-CH=CH-$.

Compounds of formula (I), in which A^2 is cycloalkyl, cycloalkyl-lower-alkyl, lower-alkenyl, lower-alkinyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy, are preferred. Further, compounds in which A^1 and A^2 are bonded to each other to form a ring and $-A^1-A^2-$ is lower-alkylene, or lower-alkenylene, optionally substituted by R^4 , in which one $-CH_2-$ group of $-A^1-A^2-$ can optionally be replaced by a NR^5 , S, or O, are preferred. Further, compounds in which A^3 and A^4 are not bonded to each other, are preferred. Compounds, in which R^3 is halogen, NH_2 , $N(lower-alkyl)_2$, $CONH_2$, CN, NO_2 , CF_3 , OH, lower-alkoxy, thio-lower-alkoxy, or is lower-alkyl, lower-alkenyl, or lower-alkinyl, optionally substituted with OH, SH or NH_2 , are also preferred.

Another preferred embodiment of the present invention relates to compounds of formula (I)



wherein

- U is O or a lone pair,
- Y N, or CR^1 ,
- Z is S, or $S(O_2)$,
- V is a) $-CH=CH-$, and m and n are 0, or
 b) $-CH_2-$, and $m+n \leq 2$, or
 c) O or NR^2 , and m is 1 to 6, n is 1 to 6, $m+n \leq 7$, or m is 1 to 3, n is 0, or
 d) S, and m is 1 to 7, n is 0 to 6, $m+n \leq 7$, or
 e) $-C \equiv C-$, and m is 0 to 7, n is 0 to 7, $m+n \leq 7$,
- Q is cycloalkyl, cycloalkyl-lower-alkyl, phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R^3 , or an alkyl, alkenyl or alkadienyl group optionally substituted by OH,
- A^1 is hydrogen, lower-alkenyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy,
- A^2 is cycloalkyl, cycloalkyl-lower-alkyl, lower-alkenyl, lower-alkinyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy, or

- A¹ and A² are bonded to each other to form a ring and -A¹-A²- is lower-alkylene, or lower-alkenylene, optionally substituted by R⁴, in which one -CH₂- group of -A¹-A²- can optionally be replaced by a NR⁵, S, or O,
- A³ and A⁴ independently from each other are hydrogen or lower-alkyl,
- 5 X is hydrogen or one or more optional halogen and/or lower-alkyl substituents,
- R¹ is hydrogen or lower-alkyl,
- R² is hydrogen or lower-alkyl,
- R³ is halogen, NH₂, N(lower-alkyl)₂, CONH₂, CN, NO₂, CF₃, OH, lower-alkoxy, thio-lower-alkoxy, or is lower-alkyl, lower-alkenyl, or lower-alkinyl,
- 10 optionally substituted with OH, SH or NH₂,
- R⁴ is hydroxy, lower-alkyl, lower-alkoxy, or thio-lower-alkoxy,
- R⁵ is hydrogen, lower-alkyl, or lower-alkyl-carbonyl,

and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

- Preferred compounds of general formula (I) are those selected from the group
- 15 consisting of
- Allyl-[2-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-methyl-amine,
 [2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-dimethyl-amine,
 Allyl-[3-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-methyl-amine,
 [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-dimethyl-amine,
- 20 Allyl-[4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-methyl-amine,
 [4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-dimethyl-amine,
 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-methyl-amino]-ethanol,
 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-methyl-amino]-ethanol,
 2-[[2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-methyl-amino]-ethanol,
- 25 [4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-(2-methoxy-ethyl)-methyl-amine,
 [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-(2-methoxy-ethyl)-methyl-amine,
 [2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-(2-methoxy-ethyl)-methyl-
- 30 amine,
 3-(4-Bromo-phenyl)-6-(4-morpholin-4-yl-butoxy)-benzo[d]isothiazole,
 3-(4-Bromo-phenyl)-6-(3-morpholin-4-yl-propoxy)-benzo[d]isothiazole,
 3-(4-Bromo-phenyl)-6-(2-morpholin-4-yl-ethoxy)-benzo[d]isothiazole,
 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-ethyl-amino]-ethanol,

- 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-ethyl-amino]-ethanol,
 2-[[2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-ethyl-amino]-ethanol,
 3-(4-Bromo-phenyl)-6-(4-pyrrolidin-1-yl-butoxy)-benzo[d]isothiazole,
 3-(4-Bromo-phenyl)-6-(3-pyrrolidin-1-yl-propoxy)-benzo[d]isothiazole,
 5 3-(4-Bromo-phenyl)-6-(2-pyrrolidin-1-yl-ethoxy)-benzo[d]isothiazole,
 6-(4-Azetidin-1-yl-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole,
 6-(3-Azetidin-1-yl-propoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole,
 6-(2-Azetidin-1-yl-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole,
 2-({4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amino)-ethanol,
 10 {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine,
 4-{4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-morpholine,
 2-({4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-ethanol,
 1-{4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-pyrrolidine,
 15 1-{4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-azetidine,
 {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-isopropyl-methyl-amine,
 {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine,
 {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
 20 Allyl-{4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amine,
 2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol,
 6-(4-Azetidin-1-yl-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole,
 3-(4-Chloro-phenyl)-6-(4-piperidin-1-yl-butoxy)-benzo[d]isothiazole,
 {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
 25 allyl-(3-[3-(4-bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-methyl-amine,
 allyl-(4-[3-(4-bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl)-methyl-amine,
 30 (3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-dimethyl-amine,
 (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl)-dimethyl-amine,
 6-(3-Azetidin-1-yl-propoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide,
 35 6-(4-Azetidin-1-yl-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide,
 {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl}-methyl-propyl-amine,
 {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-propyl-amine,

- 2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl}-ethyl-amino)-ethanol,
2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol,
5 {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl}-diethyl-amine,
{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine,
{3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl}-(2-methoxy-ethyl)-methyl-amine,
10 {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine,
2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl}-methyl-amino)-ethanol,
15 2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol,
{3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl}-ethyl-(2-methoxy-ethyl)-amine,
{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
20 {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine,
{4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-cyclopropylmethyl-methyl-amine,
{4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-isopropyl-methyl-amine,
25 3-(4-Chloro-phenyl)-6-(4-pyrrolidin-1-yl-butoxy)-benzo[d]isothiazole,
" {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine,
2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol,
(3R)-1-{4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-pyrrolidin-3-ol,
30 2-[{3-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-(2-hydroxy-ethyl)-amino]-ethanol,
{3-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-bis-(2-methoxy-ethyl)-amine,
2-[{4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-(2-hydroxy-ethyl)-amino]-ethanol,
35 {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-bis-(2-methoxy-ethyl)-amine,
2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-hydroxy-ethyl)-amino]-ethanol,

- [4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-bis-(2-methoxy-ethyl)-amine,
2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-(2-hydroxy-ethyl)-amino]-ethanol,
5 [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-bis-(2-methoxy-ethyl)-amine,
[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-ethyl-(2-methoxy-ethyl)-amine,
3-(4-Chloro-phenyl)-6-(4-piperidin-1-yl-butoxy)-benzo[d]isothiazole 1,1-dioxide,
10 Allyl-{4-[3-(4-chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amine,
6-(4-Azetidin-1-yl-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide,
{4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
15 2-({4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol,
{4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine,
2-({4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol,
20 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine,
3-(4-Chloro-phenyl)-6-(4-pyrrolidin-1-yl-butoxy)-benzo[d]isothiazole 1,1-dioxide,
1-{4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-pyrrolidin-3-ol,
25 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-isopropyl-methyl-amine,
{4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-cyclopropylmethyl-methyl-amine,
30 (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl)-bis-(2-methoxy-ethyl)-amine,
{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-dimethyl-amine N-oxide,
{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine N-oxide,
35 (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl)-bis-(2-methoxy-ethyl)-amine N-oxide,
Allyl-{3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-methyl-amine,
2-({3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-ethyl-amino)-ethanol,

- Allyl-{5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-methyl-amine,
 {5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-dimethyl-amine,
 2-({5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-methyl-amino)-
 ethanol,
- 5 {5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-(2-methoxy-ethyl)-
 methyl-amine,
 2-({5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol,
 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-bromo-phenyl)-benzo[d]isothiazole,
 Diethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine,
- 10 6-(4-Piperidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole,
 Allyl-methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine,
 6-(4-Azetidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole,
 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-
 ethanol,
- 15 2-((2-Hydroxy-ethyl)-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-
 butyl}-amino)-ethanol,
 Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine,
 2-[[3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl]-(2-hydroxy-ethyl)-
 amino]-ethanol,
- 20 Allyl-{3-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-methyl-amine,
 Allyl-{4-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine,
 {3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-dimethyl-amine,
 {4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine,
 2-({3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-methyl-amino)-ethanol,
- 25 2-({4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amino)-ethanol,
 {3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-(2-methoxy-ethyl)-methyl-
 amine,
 {4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-
 amine,
- 30 2-(Ethyl-{3-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-amino)-ethanol,
 2-(Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol,
 Ethyl-{3-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-(2-methoxy-ethyl)-
 amine,
 Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-
- 35 amine,
 2-[[3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl]-(2-hydroxy-ethyl)-
 amino]-ethanol,
 2-[[4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl]-(2-hydroxy-ethyl)-
 amino]-ethanol,

- {3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-bis-(2-methoxy-ethyl)-amine,
 {4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-bis-(2-methoxy-ethyl)-amine,
 5 1-{3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-pyrrolidine,
 1-{4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-pyrrolidine,
 1-{3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-azetidine,
 1-{4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-azetidine,
 Allyl-{4-[1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-
 10 butyl}-methyl-amine,
 2-({4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-
 butyl}-ethyl-amino)-ethanol,
 {4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-
 diethyl-amine,
 15 {4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-
 dimethyl-amine,
 6-(4-Piperidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-
 dioxide,
 6-(4-Azetidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide,
 20 2-({4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-
 butyl}-(2-hydroxy-ethyl)-amino)-ethanol,
 {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(1,1-dimethyl-prop-2-ynyl)-
 amine,
 2-{4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butylamino}-ethanol,
 25 {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-prop-2-ynyl-amine,
 {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-prop-2-ynyl-amine,
 6-(4-Azetidin-1-yl-but-1-ynyl)-3-(4-bromo-phenyl)-benzo[d]isothiazole,
 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-methyl-amino)-
 ethanol,
 30 2-({3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-methyl-amino)-
 ethanol,
 2-({3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-ethyl-amino)-ethanol,
 Allyl-{3-[3-(4-bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-propyl}-
 methyl-amine,
 35 Allyl-{4-[3-(4-bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-
 methyl-amine,
 {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-propyl}-
 dimethyl-amine,
 2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-propyl}-

- methyl-amino)-ethanol,
 2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amino)-ethanol,
 2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-propyl}-ethyl-
 5 amino)-ethanol,
 2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-ethanol,
 {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-propyl}-(2-methoxy-ethyl)-methyl-amine,
 10 {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine,
 1-{3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-propyl}-azetidine,
 1-{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-
 15 azetidine,
 Allyl-methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine,
 2-(Methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amino)-ethanol,
 20 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole,
 (2-Methoxy-ethyl)-methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine,
 Dimethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine,
 {5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-dimethyl-amine,
 25 3-(4-Chloro-phenyl)-6-(5-piperidin-1-yl-pent-1-ynyl)-benzo[d]isothiazole,
 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-chloro-phenyl)-benzo[d]isothiazole,
 {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine,
 Z-2-({3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-allyl}-ethyl-amino)-ethanol
 30 acetate,
 Allyl-{5-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-methyl-amine,
 {5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-diethyl-amine,
 2-[[5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl]-(2-hydroxy-ethyl)-amino]-ethanol,
 35 2-({5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol,
 Allyl-methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine,
 2-(Methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol,

- 2-(Ethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol,
- 6-(3-Azetidin-1-yl-prop-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole,
- (2-Methoxy-ethyl)-methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine,
- Dimethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine,
- 3-(4-Bromo-phenyl)-6-(4-piperidin-1-yl-butoxy)-benzo[d]isothiazole,
- 3-(4-Bromo-phenyl)-6-[4-(4-methyl-piperazin-1-yl)-butoxy]-benzo[d]isothiazole,
- 1-(4-(4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl)-piperazin-1-yl)-ethanone,
- Allyl-{4-[3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine,
- {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine,
- 2-({4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-ethanol,
- {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine,
- 1-{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-piperidine,
- 1-{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-azetidine,
- 2-({4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-(2-hydroxy-ethyl)-amino)-ethanol,
- {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-(2-methoxy-ethyl)-methyl-amine,
- {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-dimethyl-amine,
- Allyl-{4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-methyl-amine,
- 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-ethyl-amino)-ethanol,
- 2-(Ethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amino)-ethanol,
- 2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-ethyl-amino)-ethanol,
- {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-propyl-amine,
- {3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-dimethyl-amine,
- {3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-dimethyl-amine,
- 2-(Ethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-amino)-ethanol,
- Dimethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-amine,
- 2-(Methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-amino)-ethanol,
- Allyl-{4-[3-(4-bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine,

- {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine,
- {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-methyl-propyl-amine,
- 5 2-([4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino)-ethanol,
- 2-([4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl]-ethyl-amino)-ethanol,
- {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
- 10 1-[4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl]-azetidine,
- Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amine,
- Allyl-methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amine,
- 15 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amino)-ethanol,
- 2-(Methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amino)-ethanol,
- 20 Allyl-{4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-methyl-amine,
- {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-diethyl-amine,
- {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-methyl-propyl-amine,
- 2-([4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl]-(2-hydroxy-ethyl)-amino)-ethanol,
- 25 2-([4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl]-ethyl-amino)-ethanol,
- {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
- 6-(4-Azetidin-1-yl-butyl)-3-(4-chloro-phenyl)-benzo[d]isothiazole,
- (2-Methoxy-ethyl)-methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amine,
- 30 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-amino)-ethanol,
- Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-amine,
- 2-(Methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-amino)-ethanol,
- 35 2-([3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-propyl]-ethyl-amino)-ethanol,
- and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

Other preferred compounds of general formula (I) are those selected from the group consisting of

- 2-({5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-methyl-amino)-ethanol,
- 2-(Ethyl-{5-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amino)-ethanol,
- {5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-dimethyl-amine,
- 5 Allyl-{5-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-methyl-amine,
- Ethyl-{5-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-(2-methoxy-ethyl)-amine,
- {5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-(2-methoxy-ethyl)-methyl-amine,
- 10 2-(Ethyl-{4-[3-(4-ethynyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
- 2-(Ethyl-{4-[3-(4-ethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
- 2-((3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-1,1-dimethyl-prop-2-ynyl)-ethyl-amino)-ethanol,
- 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol N-
- 15 oxide,
- 2-(Ethyl-{4-[3-(4-piperidin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
- 2-(Ethyl-{4-[3-(4-morpholin-4-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
- 20 2-[Ethyl-(4-{3-[4-(methyl-phenyl-amino)-phenyl]-benzo[d]isothiazol-6-yloxy}-butyl)-amino]-ethanol,
- Allyl-{4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-methyl-amine,
- {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-dimethyl-amine,
- 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-methyl-amino)-ethanol,
- 25 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-amino)-ethanol,
- {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
- {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-(2-methoxy-ethyl)-methyl-amine,
- 2-{Ethyl-[4-(3-phenyl-benzo[d]isothiazol-6-yloxy)-butyl]-amino}-ethanol,
- 30 2-(Ethyl-{4-[3-(4-piperazin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
- 1-{4-[4-(6-{4-[Ethyl-(2-hydroxy-ethyl)-amino]-butoxy}-benzo[d]isothiazol-3-yl)-phenyl]-piperazin-1-yl}-ethanone,
- 2-({4-[3-(4-Benzylamino-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-
- 35 ethanol,
- 2-({4-[3-(4-Fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol,
- {4-[3-(4-Fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine,
- Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-

- amine,
 2-(Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
 2-{3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propylamino}-ethanol,
 {3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-methyl-prop-2-ynyl-
 5 amine,
 {3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-prop-2-ynyl-amine,
 3-(4-Bromo-phenyl)-6-(3-piperidin-1-yl-propoxy)-benzo[d]isothiazole,
 3-(4-Bromo-phenyl)-6-[3-(4-methyl-piperazin-1-yl)-propoxy]-benzo[d]isothiazole,
 1-(4-{3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-piperazin-1-yl)-
 10 ethanone,
 2-({4-[3-(4-Amino-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol,
 Allyl-{1,1-dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-methyl-amine,
 2-((1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl)-ethyl-amino)-ethanol,
 15 {1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-
 (2-methoxy-ethyl)-methyl-amine,
 {1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-ethyl-(2-methoxy-ethyl)-amine,
 20 2-(Ethyl-{1-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ylethynyl]-cyclohexyl}-amino)-ethanol,
 {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-fluoro-ethyl)-amine,
 Ethyl-(2-fluoro-ethyl)-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine,
 25 2-[(4-[3-[4-(3-Dimethylamino-prop-1-ynyl)-phenyl]-benzo[d]isothiazol-6-yloxy]-butyl)-ethyl-amino]-ethanol,
 3-[4-(6-{4-[Ethyl-(2-hydroxy-ethyl)-amino]-butoxy}-benzo[d]isothiazol-3-yl)-phenyl]-prop-2-yn-1-ol,
 2-[Ethyl-(4-{3-[4-(3-methylamino-prop-1-ynyl)-phenyl]-benzo[d]isothiazol-6-yloxy]-butyl)-amino]-ethanol,
 2-(Ethyl-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol,
 (2-Methoxy-ethyl)-methyl-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine,
 35 Ethyl-(2-methoxy-ethyl)-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine,
 Allyl-methyl-{1-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ylethynyl]-cyclopentyl}-amine,

- 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amine,
3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-ethyl-amine,
2-{Ethyl-[4-(3-methyl-benzo[d]isothiazol-6-yloxy)-butyl]-amino}-ethanol,
Ethyl-(2-methoxy-ethyl)-[4-(3-methyl-benzo[d]isothiazol-6-yloxy)-butyl]-amine,
5 2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-1,1-dimethyl-ethyl}-dimethyl-amine,
2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-1-methyl-ethyl}-dimethyl-amine,
1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-piperidin-4-ol,
10 4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-propyl-amine,
2-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butylamino}-ethanol,
2-(Ethyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol,
15 Allyl-methyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-amine,
1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-piperidine,
20 4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine,
2-(Ethyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol,
25 Diethyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-amine,
Methyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-propyl-amine,
Ethyl-(2-methoxy-ethyl)-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-amine,
30 2-((2-Hydroxy-ethyl)-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol,
Methyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-propyl-amine,
2-(Ethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-amino)-ethanol,
35 Diethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-amine,
Allyl-methyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-

- ynyl}-amine,
 Allyl-methyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-
 ynyl}-amine,
 Methyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yny-
 5 propyl}-amine,
 2-(Ethyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-
 ynyl}-amino)-ethanol,
 Diethyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-
 ynyl}-amine,
 10 1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-azetidine,
 1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-piperidin-
 4-ol,
 1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-piperidine,
 1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-
 15 pyrrolidine,
 2-(Ethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-
 amino)-ethanol,
 Allyl-methyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-
 amine,
 20 Allyl-methyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-
 amine,
 2-(Ethyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-
 amino)-ethanol,
 2-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butylamino}-
 25 ethanol,
 1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-piperidine,
 1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-piperidin-4-
 ol,
 {4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-propyl-amine,
 30 2-(Ethyl-{3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-
 benzo[b]thiophen-6-yl]-propyl}-amino)-ethanol,
 2-{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-
 propylamino}-ethanol,
 {3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-
 35 propyl}-propyl-amine,
 1-{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-
 propyl}-piperidin-4-ol,
 1-{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-
 propyl}-piperidine,

- Allyl-methyl-{3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-propyl}-amine,
- 2-(Ethyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl}-amino)-ethanol,
- 5 Allyl-methyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl}-amine,
- 1-{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl}-piperidine,
- 1-{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl}-piperidin-4-ol,
- 10 {4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl}-propyl-amine,
- 2-{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butylamino}-ethanol,
- 15 2-(Ethyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amino)-ethanol,
- Methyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-propyl-amine,
- Allyl-methyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amine,
- 20 Diethyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amine,
- Methyl-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine,
- 25 {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine,
- Methyl-(2-methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine,
- Methyl-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-pyrimidin-4-yl-amine,
- 30 (2-Methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine,
- {2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-pyrimidin-4-yl-amine,
- {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-(2-methyl-pyrimidin-4-yl)-amine,
- 35 {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-pyrimidin-4-yl-amine,
- {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-(2-methyl-pyrimidin-4-yl)-amine,

- {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-pyrimidin-4-yl-amine,
 {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-pyridin-4-yl-amine,
 5 1-{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-1H-imidazole, and
 1-{2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-1H-imidazole,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.
- 10 Particularly preferred compounds of general formula (I) are those selected from the group consisting of
 2-({5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol,
 2-(Ethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amino)-ethanol,
 15 2-[{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-(2-hydroxy-ethyl)-amino]-ethanol,
 {5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-dimethyl-amine,
 {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
 20 3-(4-Bromo-phenyl)-6-[4-(4-methyl-piperazin-1-yl)-butoxy]-benzo[d]isothiazole,
 {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine, and
 Allyl-methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine,
 25 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

- Other particularly preferred compounds of general formula (I) are those selected from the group consisting of
 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-ethyl-amino]-ethanol,
 Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine,
 30 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-ethyl-amino]-ethanol,
 6-(3-Azetidin-1-yl-prop-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole,
 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol,
 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol,
 35 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-cyclopropylmethyl-methyl-amine, and

Dimethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine, and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

Other particularly preferred compounds of general formula (I) are those selected from the group consisting of

- 5 {5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-diethyl-amine,
2-({5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol,
Allyl-{4-[3-(4-bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-
butyl}-methyl-amine,
{4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-
10 propyl-amine,
2-[[4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-
butyl]-(2-hydroxy-ethyl)-amino]-ethanol,
6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole, and
2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amino)-
15 ethanol,
and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

Other particularly preferred compounds of general formula (I) are those selected from the group consisting of

- 20 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-
methoxy-ethyl)-amine,
{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-dimethyl-
amine, and
2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol,
and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

Other particularly preferred compounds of general formula (I) are those selected from the group consisting of

- 3-(4-Bromo-phenyl)-6-(3-piperidin-1-yl-propoxy)-benzo[d]isothiazole,
2-(Ethyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-
amino)-ethanol,
25 2-(Ethyl-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-
ynyl}-amino)-ethanol,
Ethyl-(2-methoxy-ethyl)-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-
benzo[b]thiophen-6-yloxy]-butyl}-amine,
{4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
35 2-(Ethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-
amino)-ethanol,

- 2-((1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl)-ethyl-amino)-ethanol,
 2-((3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-1,1-dimethyl-prop-2-ynyl)-ethyl-amino)-ethanol,
 5 Methyl-(2-methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine, and
 (2-Methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.


- 10 Compounds of formula (I) can have one or more asymmetric carbon atoms and can exist in the form of optically pure enantiomers or as racemats. The invention embraces all of these forms.

It will be appreciated, that the compounds of general formula (I) in this invention may be derivatised at functional groups to provide derivatives which are capable of

- 15 conversion back to the parent compound in vivo.


The present invention also relates to a process for the manufacture of compounds as described above, which process comprises

- a) reacting a compound of formula (II)




- 20 with a compound $(A^1, A^2, U)N-C(A^3, A^4)-(CH_2)_m-M$, wherein V is O, S or NR^2 , M is mesylate, tosylate, triflate, Cl, Br or I, and X, Y, Z, Q, U, A^1 , A^2 , A^3 , A^4 , m, n and R^2 have the significances given above, or wherein HV is mesylate, tosylate, triflate, Cl, Br or I, and M is OH, SH or NHR^2 , and R^2 has the significance given above,

- or b) reacting a compound of formula (III)




with a compound NHA^1A^2 , wherein M is mesylate, tosylate, triflate, Cl, Br or I, and $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4, \text{V}, \text{X}, \text{Y}, \text{Z}, \text{Q}, \text{m}$ and n are as defined above,

or c) reacting a compound of formula (IV)




5 with a compound $(\text{A}^1, \text{A}^2, \text{U})\text{N}-\text{C}(\text{A}^3, \text{A}^4)-(\text{CH}_2)_m-\text{C}\equiv\text{CH}$, wherein $\text{X}, \text{Y}, \text{Z}, \text{Q}, \text{U}, \text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4$ and m are as defined above,

or d) reacting a compound of formula (V)



10 with a compound $(\text{A}^1, \text{A}^2, \text{U})\text{N}-\text{C}(\text{A}^3, \text{A}^4)-(\text{CH}_2)_m-\text{M}$, wherein M is mesylate, tosylate, Cl, Br or I, and $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4, \text{U}, \text{X}, \text{Y}, \text{Z}, \text{Q}, \text{m}$ and n are as defined above,

or e) hydrogenating a compound of formula (VI)



wherein V is $-\text{C}\equiv\text{C}-$, and $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4, \text{U}, \text{X}, \text{Y}, \text{Z}, \text{Q}, \text{m}$ and n are as defined above,

and optionally converting a compound of formula (I) as defined above, wherein U is a
15 lone pair, to a corresponding compound wherein U is O.

The invention further relates to compounds of formula (I) as defined above, when manufactured according to a process as defined above.

As described above, the compounds of formula (I) of the present invention can be used for the treatment and/or prophylaxis of diseases which are associated with OSC such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections and gallstones, and/or treatment and/or prophylaxis of impaired glucose tolerance, diabetes, tumors and/or hyperproliferative disorders, preferably for the treatment and/or prophylaxis of hypercholesterolemia and/or hyperlipemia. Hyperproliferative skin and vascular disorders particularly come into consideration as hyperproliferative disorders.

The invention therefore also relates to pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

Further, the invention relates to compounds as defined above for use as therapeutic active substances, particularly as therapeutic active substances for the treatment and/or prophylaxis of diseases which are associated with OSC such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones, tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance and diabetes, preferably for the treatment and/or prophylaxis of hypercholesterolemia and/or hyperlipemia.

In another embodiment, the invention relates to a method for the treatment and/or prophylaxis of diseases which are associated with OSC such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones, tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance and diabetes, preferably for the treatment and/or prophylaxis of hypercholesterolemia and/or hyperlipemia, which method comprises administering a compound as defined above to a human being or animal.


The invention further relates to the use of compounds as defined above for the treatment and/or prophylaxis of diseases which are associated with OSC such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones, tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance and diabetes, preferably for the treatment and/or prophylaxis of hypercholesterolemia and/or hyperlipemia.

In addition, the invention relates to the use of compounds as defined above for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with OSC such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones, tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance and

diabetes, preferably for the treatment and/or prophylaxis of hypercholesterolemia and/or hyperlipemia. Such medicaments comprise a compound as defined above.


The compounds of formula (I) can be manufactured by the methods given below, by the methods given in the examples or by analogous methods. Appropriate reaction
5 conditions for the individual reaction steps are known to the person skilled in the art. Starting materials are either commercially available or can be prepared by methods analogous to the methods given in the examples or by methods known in the art.

Scheme 1




PG = protecting group

Scheme 2




PG = protecting group

Scheme 3




PG = protecting group, such as trityl for sulfur or BOC for NR²

Scheme 4



Scheme 6



Scheme 1:

For compounds which are benzo[d]isothiazole derivatives, the synthesis starts from a suitable protected fluoro-substituted aromatic system 1 ($V = O$ or NR^2 , and $n = 0-6$, or $V=S$ and $n=0$) and an acid chloride 2 which is converted under Friedel-Crafts conditions with a Lewis acid such as aluminium chloride to the phenone 3 (step a). The starting materials 1 are either commercially available or can be obtained by methods known in the art, e.g. from the corresponding acids. The transformation to the heterocyclic system may be achieved by reacting the fluoride 3 with e.g. benzylmercaptane in the presence of potassium tert.butyrate in THF at RT, treating the benzylthio ether 4 obtained with sulfonyl chloride in methylene chloride (at 0 °C) and cyclizing the resulting intermediate in THF with a solution of ammonia in ethanol to 5, as described in European Patent application 778271 (steps b and c). In case of methoxyphenylether 5 ($n=0$, $V-PG=OMe$) treatment with boron tribromide in methylene chloride at -78°C to RT or with 62% aqueous HBr in acetic acid at reflux gives phenol 6. For $V=NR^2$ or $V=O$ and $n>0$, deprotection using procedures known in the art (step d) gives the free HV-building block 6.

Oxidation of a suitable protected benzo[d]isothiazols 5 ($V = O$ or NR^2 , and $n = 0-6$, or $V=S$ and $n=0$) may be effected with e.g. $KMnO_4$ on silicagel in CH_2Cl_2 at 50 °C to yield the corresponding benzo[d]isothiazole 1,1-dioxide.

For $V=S$ and $n>0$, the sulfur is introduced in a later step (see Scheme 4).

Scheme 2:

For compounds which are benzo[b]thiophene derivatives, the synthesis starts from a suitable protected thiol 1 and α -bromoacetophenone derivative 2 which are converted to the thioether 3 by treatment with KOH in EtOH at 0°C to RT (step a). The starting materials 1 are either commercially available or can be obtained by methods known in the art, e.g. from the corresponding acids. The cyclization can be carried out by heating the compound 3 in polyphosphoric acid to 80-100°C or by treatment with $BF_3 \cdot Et_2O$ to give benzo[b]thiophene 4 (step b).

In the case of methoxyphenylether 4 ($n=0$, $V-PG=OMe$), treatment with boron tribromide in methylene chloride at -78°C to RT or with 62% aqueous HBr in acetic acid at reflux gives phenol 5. For $V=S$, NR^2 or $V=O$ and $n>0$, deprotection using procedures known in the art (step d) gives the free HV-building block 5.

Oxidation of a suitable protected benzo[b]thiophene 4 ($V = O$ or NR^2 , and $n = 0-6$, or $V=S$ and $n=0$) may be effected with e.g. sodium perborate in acetic acid at $50^\circ C$ yield the corresponding benzo[b]thiophene 1,1-dioxide 4.

For $V=S$ and $n>0$, the sulfur is introduced in a later step (see Scheme 4).

- 5 Alternatively compound of the formula 5 may be prepared as described previously in European Patent application 778271.

Scheme 3:

Scheme 3 shows the synthesis of amino-VH sidechain 4 that may be used for the synthesis of compounds with the corresponding V-spacers ($V = NR^2$, S, or O). α,ω -
10 dihaloalkane or mesyl-alkanyl-halogenide 1 may be treated with a suitable protected amine (HNR^2-PG , PG =protecting group, e.g. BOC) in DMA or a thiol ($HS-PG$) e.g., triphenylmethanethiol in the presence of NaH in DMA to give the compound 2 (step a). Treatment with the amine A^1A^2NH yields the S- or N-protected amine 3 (step b) and in the case of α,ω -haloalkanol 1 directly amino-alcohol 4. N-deprotection with procedures
15 known in the art e.g. TFA in CH_2Cl_2 gives the amine side chain 4 (step c). The deprotection of the thiol moiety in 3 may be achieved with TFA/triisopropylsilane in CH_2Cl_2 at $0^\circ C$ to RT to yield the aminothiols 4 (step c). Aminoalkanol 4 can further be transformed to mesylate 5 (step d).

Scheme 4:

- 20 Alkylation of the phenol 1 ($V=O$, S, $n=0$) in acetone with K_2CO_3 and a suitable dihaloalkane (halogene is here represented by bromine, but can also be chlorine or iodine. It is also possible to use mesylates or tosylates instead of halogenides) at reflux yields halogenide 2 (reaction step a). For the preparation of derivatives 2 ($V=O$, $n>0$), the alcohol 1 can be treated with α,ω -dihaloalkanes under phase transfer conditions e.g. α,ω -
25 dihaloalkanes, NaOH, nBu_4NHSO_4 . For $V=S$, O or NR^2-PG , the derivative 1 may be treated with α,ω -dihaloalkane in the presence of NaH in DMF $0^\circ C$ to RT to yield bromide 2. For shorter alkanes ($m=1$ or 2), the method of choice is the in situ generation of the haloalkane-triflate (from the corresponding haloalkanol with trifluoromethanesulfonic anhydride/2,6-di-tert-butylpyridine in CH_2Cl_2 at $0^\circ C$). This haloalkane-triflate may then
30 be reacted with 1 in the presence of a base such as with 2,6-di-tert-butylpyridine as base in nitromethane at $60^\circ C$ to yield bromide 2 [analogously to a procedure of Belostotskii, Anatoly M.; Hassner, Alfred. Synthetic methods. 41. Etherification of hydroxysteroids via

triflates. Tetrahedron Lett. (1994), 35(28), 5075]. Compound 2 can be converted (reaction step b) to the amine 3 with an excess of the corresponding amine NHA^1A^2 in a suitable solvent such as DMA, DMF, MeOH at RT or 50-65°C or with amine NHA^1A^2 , sodium hydride in DMF, DMA or THF. Alternatively, the compound 1 may be transferred to the amine 3 by attaching the pre-assembled fragment $\text{A}^1\text{A}^2\text{NC}(\text{A}^3\text{A}^4)(\text{CH}_2)_m$ -OMes/halogenide, which can be synthesised by known methods (shown e.g. in Scheme 3), using alkylating conditions (step d). Heteroaromate 1 ($\text{V}=\text{O}$, $n>0$) can also be mesylated 1 ($\text{V}=\text{OMes}$) and then reacted with $\text{A}^1\text{A}^2\text{NC}(\text{A}^3\text{A}^4)(\text{CH}_2)_m\text{-VH}$ (synthesis described Scheme 3) in e.g. DMF with NaH as base to give 3 (with $\text{V}=\text{O}$, S, NR^2).

10 Amine 3 may be converted to a salt or to the N-oxide 4 (step c). For the formation of N-oxide 4 ($\text{V}=\text{O}$) a mixture of hydrogen peroxide urea adduct and phthalic anhydride in CH_2Cl_2 at RT may be used. For the preparation of the N-oxides 4 ($\text{V}=\text{S}$ or NR^2) step e may be performed: Oxidation of the pre-assembled fragment $\text{A}^1\text{A}^2\text{NC}(\text{A}^3\text{A}^4)(\text{CH}_2)_m$ -OMes/halogenide to the corresponding N-oxide derivative, followed by alkylation of the compound 1.

Oxidation of 2 ($\text{V}=\text{O}$ or NR^2) may be effected with e.g. KMnO_4 on silicagel in CH_2Cl_2 at 50 °C for benzo[d]isothiazols to give the corresponding benzo[d]isothiazole 1,1-dioxide 5. In the case of benzo[b]thiophene 2, oxidation with sodium perborate in acetic acid at 50°C gives benzo[b]thiophene 1,1-dioxide 5. Bromide 5 can be transformed to 3 and 4 as described for bromide 2.

If $\text{A}^2 = \text{H}$, heteroaromatic moieties A^2 may be introduced by treatment with halo heteroaromatics in the presence of Huenig's base in DMF (Ger. Offen. (1990), DE3905364 A1). Alternatively, Buchwald conditions e.g. $\text{Pd}(\text{OAc})_2$, 2-(Dicyclohexylphosphino) biphenyl, NaOtBu in toluene might be applied (John P. Wolfe, Hiroshi Tomori, Joseph P. Sadighi, Jingjun Yin, and Stephen L. Buchwald, J. Org. Chem., 65 (4), 1158 -1174, 2000).

Amine 3 may be further modified. For Q = methyl, deprotonation with e.g. lithium diisopropylamide in THF at -78°C and then treatment with a bromide $\text{BrCH}_2\text{-R}^7$ or aldehyde $\text{HC}(\text{O})\text{-R}^7$, wherein R^7 is an alkyl, alkenyl or alkadienyl group with up to 12 C atoms, may yield the corresponding compound 3 in which Q stands for $\text{CH}_2\text{CH}_2\text{-R}^7$ or $\text{CH}_2\text{CH}(\text{OH})\text{-R}^7$.

For an amine 3 or also starting compound 1 in which Q is a halo- or hydroxy-substituted aromatic system (in case of the later, the corresponding triflate may be synthesized) the corresponding alkyne, alkyl, alkene, amine, alkoxy or thioalkoxy

substituted derivative can be synthesized employing Sonogashira reaction or palladium catalyzed amination, C-O or C-S coupling reactions. For the Sonogashira reaction of the arylhalogenide or aryltriflate may be treated with a suitable alkynes or alkynol in THF in the presence of a base such as triethylamine or piperidine with a catalytic amount of e.g.

5 Pd(PPh₃)₄/ CuI or Pd(OAc)₂/CuI or PdCl₂(PPh₃)₂/CuI at 45°C to 80°. These alkynes can then selectively be reduced. The introduction of an amine moiety may be achieved using primary or secondary amines and the arylhalogenide or aryltriflate using methods developed by Buchwald e.g. tris(Dibenzylideneacetone)dipalladium, 2(di-tertbutylphosphino)Biphenyl in toluene and Natrium tert-butylat as base to give the newly

10 substituted 3 [e.g. Wolfe, John P.; Tomori, Hiroshi; Sadighi, Joseph P.; Yin, Jingjun; Buchwald, Stephen L. Simple, efficient catalyst system for the palladium-catalyzed amination of aryl chlorides, bromides, and triflates. J. Org. Chem. (2000), 65(4), 1158-1174].

For the oxygen analogues see for example Palucki, Michael; Wolfe, John P.;

15 Buchwald, Stephen L. Palladium-Catalyzed Intermolecular Carbon-Oxygen Bond Formation: A New Synthesis of Aryl Ethers. J. Am. Chem. Soc. (1997), 119(14), 3395-3396.

These newly substituted amines 3 can then again be transformed to compound 4. For the preparation of the N-oxides 4 (V=S or NR²) step e may be performed: Oxidation of the pre-assembled fragment A¹A²NC(A³A⁴)(CH₂)_m-OMes/halogenide to the

20 corresponding N-oxide derivative, followed by alkylation of the compound 1.

Scheme 5:

In Scheme 5, the preparation of compounds of formula 6, in which V represents -CH₂-, -CH=CH- or -C≡C- is outlined starting from hydroxyphenyl derivative 1, which may be transformed to the triflate 2 in pyridine with trifluoromethanesulfonic anhydride

25 at 0 °C to RT (reaction step a). Sonogashira-coupling (reaction step b) of the triflate 2 and a suitable alkynol or alkynchloride in piperidine with Pd(PPh₃)₄/CuI at 45 °C to 80 °C in analogy to a literature procedure yields alcohol 3a or chloride 3b [Stara, Irena G.; Stary, Ivo; Kollarovic, Adrian; Teply, Filip; Saman, David; Fiedler, Pavel. Coupling reactions of halobenzenes with alkynes. The synthesis of phenylacetylenes and symmetrical or

30 unsymmetrical 1,2-diphenylacetylenes. Collect. Czech. Chem. Commun. (1999), 64(4), 649-672.]. In case of Q = Bromo or Iodo-substituted aromatic system, triflate 2 is dissolved in THF with PdCl₂(PPh₃)₂ as catalyst and alkynol or alkynchloride, triphenylphosphine, triethylamine and a catalytic amount of CuI to give alkyne 3a or 3b. Mesylation for alcohol 3a with methanesulfonylchloride e.g. in pyridine with DMAP

(reaction step c) and subsequent amination (reaction step d) of the resulting mesylate 4 with a suitable amine NHA^1A^2 in DMA at RT or as described in Scheme 4 yields the amine 5. Alcohol 3a can also be treated with trifluoromethane sulfonic acid anhydride and Huenig's base at -15°C in CH_2Cl_2 (in situ generation of the corresponding triflate) followed by treatment with the corresponding amine NHA^1A^2 at -15°C to RT. This is especially the method of choice for but-3-yn-1-ol-derivatives 3a. Chloride 3b can be transformed directly or via iodide (Finkelstein reaction) to the amine 5, as described above (step d). Compounds 5 in which V is $-\text{CH}_2-$ or $-\text{CH}=\text{CH}-$ can be obtained by hydrogenation of compound 5 in EtOH with $\text{Pt}_2\text{O} \cdot \text{H}_2\text{O}$ (yields the saturated analogue 5) or by selective hydrogenation with other known methods (yields the double bond analogue 5). Optionally, the hydrogenation described above can be performed at an earlier stage e.g. the alcohol 3a or mesylate 4.

For compounds in which A^3 and/or A^4 are not H, the group $\text{A}^1\text{A}^2\text{NC}(\text{A}^3\text{A}^4\text{C})[\text{CH}_2]_m\text{-acetylene}$ can be synthesised by known methods and attached to compound 2 (Sonogashira-coupling), to yield the compounds of the present invention 5 (reaction step f).

Compounds of the formula 5 ($n>0$) may be synthesised by Swern oxidation of the alcohol 1 ($\text{V}=\text{O}$ and $n>0$) to the corresponding aldehyde 7 (step g). The aldehyde 7 may be treated with triphenylphosphine, tetra-bromo-methane and triethylamine in CH_2Cl_2 at 0°C to RT to yield 2,2-Dibromo-vinyl derivative 8 (step h). Rearrangement with $n\text{-BuLi}$ (ca 1.6 M in hexane) in THF at -78°C , followed by reaction with formaldehyde (-78°C to RT) leads to the propargyl alcohol 9a (step i, side chain extension through application of the Corey-Fuchs method), following conditions described in: Marshall, James A.; Bartley, Gary S.; Wallace, Eli M. Total Synthesis of the Pseudopterane (-)-Kallolide B, the Enantiomer of Natural (+)-Kallolide B. J. Org. Chem. (1996), 61(17), 5729-5735; and Baker, Raymond; Boyes, Alastair L.; Swain, Christopher J. Synthesis of talaromycins A, B, C, and E. J. Chem. Soc., Perkin Trans. 1 (1990), (5), 1415-21.

For longer side chains, the rearrangement is performed with $n\text{-BuLi}$ (ca 1.6 M in hexane) in THF at -78°C as above, followed by addition of a cosolvent such as DMPU and reaction with O-protected 1-bromo-alcohols or α -chloro- ω -iodoalkane (e.g. 1-bromo- n -tetrahydro-pyran-2-ylmethoxyalkane) to yield the O-protected compounds 9b or the chloro compound 9c (step i). O-protected compounds 9b can be deprotected to the corresponding alkynol 9a (in MeOH at $50\text{-}60^\circ\text{C}$, in the presence of catalytic amount of pyridinium toluene-4-sulfonate). Alcohol 9a can be reacted with Huenig's

base/trifluoromethane sulfonic acid anhydride at -15°C in CH_2Cl_2 (in situ generation of the corresponding triflate) followed by treatment with Huenig's base and the corresponding amine NHA^1A^2 at -15°C to RT to give amine 5. Alternatively, mesylation of alcohol 9a with methanesulfonylchloride, pyridine and DMAP in CH_2Cl_2 at 0°C to RT gives mesylate 10. Conversion of the mesylate 10 or the chloride 9c (or the in situ generated iodide) to the amine 5 can be accomplished with an excess of the corresponding amine NHA^1A^2 in DMA at RT or as described above (step I).

Optionally compounds 5 in which V is $-\text{CH}_2-$ or $-\text{CH}=\text{CH}-$ can be obtained by hydrogenation of compound 5 itself or the intermediates 9a, 9b, 9c or 10. The hydrogenation may be done in EtOH with $\text{Pt}_2\text{O} \cdot \text{H}_2\text{O}$ (yields the saturated analogues 5, 9a, 9b, 9c or 10) or by selective hydrogenation to the double bond with other known methods and transforming the intermediates afterwards to 5.

For the introduction of the group $\text{A}^1\text{A}^2\text{N}(\text{A}^3\text{A}^4\text{C})[\text{CH}_2]_m$ in compound 8 to obtain the final compound 5 in which A^3 and/or A^4 are not H, the following steps have to be performed (step m or step i and I): for $m > 0$, the building block $\text{A}^1\text{A}^2\text{N}(\text{A}^3\text{A}^4\text{C})[\text{CH}_2]_m$ -halogenide/mesylate is synthesised by known methods (or in analogy to the methods described in Scheme 3) and introduced (step m) under the same condition as described above for step i. For $m = 0$, the introduction of the group $\text{A}^1\text{A}^2\text{N}(\text{A}^3\text{A}^4\text{C})$ with A^3 and/or A^4 not H, a two step procedure has to be followed: first the rearrangement of 8 with $n\text{-BuLi}$ (ca 1.6 M in hexane) in THF at -78°C , followed by reaction with the corresponding aldehyde (A^3 or $\text{A}^4\text{-COH}$) or ketone (A^3COA^4 , at -78°C to RT) leads to the A^3A^4 substituted propargyl alcohol 9a (step i) which is mesylated and reacted with the desired A^1A^2 -substituted-amine to yield A^3A^4 -substituted compound 5 (step I).

If $\text{A}^2 = \text{H}$, heteroaromatic moieties A^2 may be introduced by treatment with halo heteroaromatics in the presence of Huenig's base in DMF (Ger. Offen. (1990), DE3905364 A1). Alternatively, Buchwald conditions e.g. $\text{Pd}(\text{OAc})_2$, 2-(Dicyclohexylphosphino) biphenyl, NaOtBu in toluene might be applied (John P. Wolfe, Hiroshi Tomori, Joseph P. Sadighi, Jingjun Yin, and Stephen L. Buchwald, J. Org. Chem., 65 (4), 1158 -1174, 2000).

Amine 5 can further be transformed. For $\text{Q} = \text{methyl}$ as described in Scheme 4 to the corresponding compound 5 in which Q stands for $\text{CH}_2\text{CH}_2\text{-R}^7$ or $\text{CH}_2\text{CH}(\text{OH})\text{-R}^7$.

For an amine 5 in which Q is a halo-substituted aromatic system, the corresponding alkyne, alkyl, alkene, amine, alkoxy or thioalkoxy substituted derivative can be synthesized employing Sonogashira reaction or palladium catalyzed amination, C-O or C-S coupling

reactions (as described for compound 3 in scheme 4) (or in case of a hydroxy substitution, the corresponding triflate may be synthesized).

Amine 5 may be converted to a salt or to the N-oxide 6 using a mixture of hydrogen peroxide urea adduct and phthalic anhydride in CH_2Cl_2 at RT (step e).

5 Scheme 6:

Another approach for the introduction of the substituted side chain is depicted in scheme 6. It starts by alkylation of the phenol 1 ($\text{V}=\text{O}$, S, $n=0$) directly with a suitable ω -halo-alkane carboxylic ester or via the in situ generated triflate of an ω -hydroxy alkane carboxylic ester in analogy to Belostotskii, Anatoly M.; Hassner, Alfred. Synthetic
10 methods. 41. Etherification of hydroxysteroids via triflates. Tetrahedron Lett. (1994), 35(28), 5075-6(step a). For the preparation of derivatives 2 ($\text{V}=\text{O}$, $n>0$), the alcohol 1 can be treated with ω -haloalkane carboxylic ester (halogene is here represented by bromine, but can also be chlorine or iodine. It is also possible to use mesylates or tosylates or triflates instead of halogenides) under phase transfer conditions. For $\text{V}=\text{S}$, O or NR^2 -PG, the
15 derivative 1 may be treated with ω -haloalkane carboxylic ester in the presence of NaH in DMF 0 °C to RT to yield ester 2.

Alternatively, the ester 2 can be prepared from the bromide 3 (synthesis according to scheme 4) by treatment with e.g. acetocyanhydride in acetonitrile, followed by a Pinner reaction and hydrolysis of the imidate to the corresponding ester (step b).

20 For $\text{V}=\text{CH}=\text{CH}$, the ester 2 or its corresponding acid may be prepared from aldehyde 4 (synthesis described in scheme 5) by treatment with the corresponding Wittig reagent $\text{XPh}_3\text{P}^+(\text{CH}_2)_{m+1}\text{CO}_2\text{R}/\text{H}$. For $\text{V}=\text{C}$, hydrogenation of the Wittig product under standard conditions yields the saturated product 2.

For $\text{V}=\text{C}\equiv\text{C}$, ester 2 or amide 6 may be derived from the dibromoderivative 5
25 (synthesis according to scheme 5) by rearrangement with $n\text{-BuLi}$ (ca 1.6 M in hexane) in THF at -78°C , followed by reaction with chloroformate or dimethylcarbamoyl chloride (-78°C to RT; step d). For longer side chains, the rearrangement of dibromoalkene 5 may be performed with $n\text{-BuLi}$ (ca 1.6 M in hexane) in THF at -78°C as above, followed by addition of a cosolvens such as DMPU and reaction with a suitable protected 1-bromo-
30 alkylalcohol $\text{Br}-(\text{CH}_2)_m\text{CH}_2\text{OH}$, followed by oxidation to yield the compound 2 as acid (step e).

Saponification of the ester 2 using standard conditions e.g. LiOH in EtOH, MeOH or THF, followed by treatment with NHA^1A^2 , EDCI, HOBT and a base such as Huenig's base, NEt_3 , NMM in CH_2Cl_2 , DMF, DMA or dioxane gives amide 6. Amide 6 can be transferred to amine 7 ($\text{A}^3, \text{A}^4 = \text{Me}$) by reaction with methylmagnesium bromide, ZrCl_4 in THF at low temperature (see Stephen M. Denton, Anthony Wood, A Modified Bouveault Reaction for the Preparation of α, α -dimethylamines from Amides, *Synlett* 1999, 1, 55-56.) or by treatment with other grignard reagents in the presence of ZrCl_4 or $\text{Ti}(\text{OiPr})_4$ (see V. Chalinski, A. de Meijere, A versatile New Preparation of Cyclopropylamines from acid dialkylamides, *Angew. Chem. Int. Ed. Engl.* 1996, 35, No4, 413-4.).

10 For $\text{A}^1 = \text{Me}$, $\text{A}^2 = \text{OMe}$, the amide 3 can be treated with a grignard reagent A^3MgX to give the corresponding ketone 8. Reductive alkylation of the ketone 8 by treatment with NHA^1A^2 in the presence of tetraisopropyl orthotitanate, followed by reduction with NaCNBH_3 in ethanol yields the amine 7 (see: R. J. Mattson, K. M. Pham, D. J. Leuck, K.A. Cowen, *J.O.C.* 1990, 55, 2552-4.).

15 Amine 7 may be converted to a salt or to the N-oxide 9 using a mixture of hydrogen peroxide urea adduct and phthalic anhydride in CH_2Cl_2 at RT.

Amine 7 can be transformed further. For Q = methyl as described in Scheme 4 to the corresponding compound 7 in which Q stands for $\text{CH}_2\text{CH}_2\text{-R}^7$ or $\text{CH}_2\text{CH}(\text{OH})\text{-R}^7$.

20 For an amine 7 in which Q is a halo-substituted aromatic system, the corresponding alkyne, alkyl, alkene, amine, alkoxy or thioalkoxy substituted derivative can be synthesized employing Sonogashira reaction or palladium catalyzed amination, C-O or C-S coupling reactions (as described for compound 3 in scheme 4) (or in case of a hydroxy substitution, the corresponding triflate may be synthesized).

The following tests were carried out in order to determine the activity of the compounds of formula I and their salts.

Inhibition of human liver microsomal 2,3-oxidosqualene-lanosterol cyclase (OSC)

Liver microsomes from a healthy volunteer were prepared in sodium phosphate
5 buffer (pH 7.4). The OSC activity was measured in the same buffer, which also contained
1mM EDTA and 1mM dithiothreitol. The microsomes were diluted to 0.8mg/ml protein
in cold phosphate buffer. Dry [^{14}C]R,S-monooxidosqualene (MOS, 12.8 mCi/mmol) was
diluted to 20 nCi/ μl with ethanol and mixed with phosphate buffer-1% BSA (bovine
serum albumin). A stock solution of 1 mM test substance in DMSO was diluted to the
10 desired concentration with phosphate buffer-1% BSA. 40 μl of microsomes were mixed
with 20 μl of the solution of the test substance and the reaction was subsequently started
with 20 μl of the [^{14}C]R,S-MOS solution. The final conditions were: 0.4mg/ml of
microsomal proteins and 30 μl of [^{14}C]R,S-MOS in phosphate buffer, pH 7.4, containing
0.5% albumin, DMSO <0.1% and ethanol <2%, in a total volume of 80 μl .

15 After 1 hour at 37°C the reaction was stopped by the addition of 0.6 ml of 10%
KOH-methanol, 0.7ml of water and 0.1ml of hexane:ether (1:1, v/v) which contained 25 μg
of non-radioactive MOS and 25 μg of lanosterol as carriers. After shaking, 1 ml of
hexane:ether (1:1, v/v) was added to each test tube, these were again shaken and then
centrifuged. The upper phase was transferred into a glass test tube, the lower phase was
20 again extracted with hexane:ether and combined with the first extract. The entire extract
was evaporated to dryness with nitrogen, the residue was suspended in 50 μl of
hexane:ether and applied to a silica gel plate. Chromatographic separation was effected in
hexane:ether (1:1, v/v) as the eluent. The R_f values for the MOS substrate and the
lanosterol product were 0.91 and, respectively, 0.54. After drying, radioactive MOS and
25 lanosterol were observed on the silica gel plate. The ratio of MOS to lanosterol was
determined from the radioactive bands in order to determine the yield of the reaction and
OSC inhibition.

The test was carried out on the one hand with a constant test substance
concentration of 100nM and the percentage OSC inhibition against controls was
30 calculated. The more preferred compounds of the present invention exhibit inhibitions
larger than 50%. In addition, the test was carried out with different test substance
concentrations and subsequently the IC₅₀ value was calculated, i.e. the concentration
required to reduce the conversion of MOS into lanosterol to 50% of the control value. The
preferred compounds of the present invention exhibit IC₅₀ values of 1 nM to 10 μM ,
35 preferably of 1 - 100 nM.

The compounds of formula I and their pharmaceutically acceptable acid addition salts can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories, parenterally, e.g. in the form of injection solutions or infusion solutions, or topically, e.g. in the form of ointments, creams or oils.

The production of the pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described compounds of formula I and their pharmaceutically acceptable acid addition salts, optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers are, however, required in the case of soft gelatine capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

The dosage of the compounds of formula I can vary within wide limits depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 mg to about 1000 mg, especially about 50 mg to about 500 mg, comes into consideration for the prevention and

control of topical and systemic infections by pathogenic fungi. For cholesterol lowering and treatment of impaired glucose tolerance and diabetes the daily dosage conveniently amounts to between 1 and 1000mg, preferably 5 to 200mg, for adult patients. Depending on the dosage it is convenient to administer the daily dosage in several dosage units.

- 5 The pharmaceutical preparations conveniently contain about 1-500 mg, preferably 5-200 mg, of a compound of formula I.

The following Examples serve to illustrate the present invention in more detail. They are, however, not intended to limit its scope in any manner.

Examples

Abbreviations:

- AcOH = acetic acid, Aq = aqueous, BOC = t-butyloxycarbonyl, CH₂Cl₂ = dichloromethane, DAST = Diethylaminosulfur trifluoride, EDCI = N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride, Et₃N = triethylamine, Et₂O = diethylether, EtOAc = ethylacetate, EtOH = ethanol, eq = equivalents, HOBT = 1-Hydroxybenzo-triazole, Huenigs base = iPr₂NEt N-ethyl-diisopropylamine, LAH = Lithium aluminium hydride, LiBH₄ = lithium borohydride, MeOH = methanol, NMM = N-methyl-morpholine, TBDMSCl = t-butyldimethylsilyl chloride, Pd(Ph₃P)₄ = tetrakis(triphenylphosphine)palladium, PdCl₂(PPh₃)₂ = bis(triphenylphosphine)palladium(II) chloride, TFA = trifluoroacetic acid, DMAP = 4-Dimethylaminopyridine, THF = tetrahydrofuran.

General remarks

All reactions were performed under argon.

- 15 The purification of the final amines by preparative HPLC [e.g. RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile] yielded mixtures of the corresponding amino formate and the corresponding halogenide or mesylate which was used in the reaction. The ratio was not always determined, the purity of the final amino salts was >80% after LC-MS.

Example 1

1.1

1.4 l of nitrobenzene were cooled in an ice bath and treated in succession at a maximum temperature of 6°C with 492.6 g (3.656 mol) of aluminum chloride and 700 g (3.355 mol) of 4-trifluoromethyl-2-fluoro-benzoyl chloride in 350 ml of nitrobenzene. The mixture was stirred (15 min) and 427.5 g (3.388 mol) of 3-fluoroanisole were added slowly so that the temperature did not rise above 6°C. The solution was left to warm to room temperature, stirred 2h at this temperature, poured on to ice-water (2.5 l) and extracted with methylene chloride. The organic phase was washed with water and concentrated. After crystallization from THF (1.2l)/hexane (4.2l), 237 g (24%) of (2-Fluoro-4-methoxy-phenyl)-(4-trifluoromethyl-phenyl)-methanone were obtained, MS: 298 (M).

1.2

12.01 g (107 mmol) of potassium tert. butylate were dissolved in 425 ml of THF and treated slowly with 13.29 g (107 mmol) of benzyl mercaptane. The suspension was stirred at RT (1 h) and then treated with 29.56 g (99 mmol) of (2-Fluoro-4-methoxy-phenyl)-(4-trifluoromethyl-phenyl)-methanone in 220 ml of THF. The solution was stirred at RT (2 h), evaporated and treated with EtOAc/ aqueous NaHCO₃ solution and dried (Na₂SO₄). 42.8 g (corresponds to 99 mmol) of (2-Benzylsulfanyl-4-methoxy-phenyl)-(4-trifluoromethyl-phenyl)-methanone were obtained which were taken up in 340 ml of methylene chloride, treated at 0 °C with 13.38 g (99 mmol) of sulphuryl chloride and stirred at 0 °C for 90 min. After evaporation, the residue was taken up in 307 ml of THF, treated at 0 °C with 307 ml of a saturated ammonia solution in ethanol and stirred (16 h at RT). The solvent was removed and the residue was again taken up in sodium hydrogen carbonate solution and ethyl acetate. The phases were separated and the inorganic phase was extracted with ethyl acetate. The organic phases were washed with sodium chloride solution and dried. Recrystallization from CH₂Cl₂/methanol (-20 °C) yielded 19.8 g (71%) of 6-Methoxy-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole, MS: 309 (M).

1.3

A solution of 20 g (64.66 mmol) of 6-Methoxy-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole in 150 ml of acetic acid and 75 ml 62% aqueous HBr solution was heated to reflux for 70 h, subsequently cooled and taken up in ice water and extracted with tert-butylmethyleter (3x). The organic phases were washed with saturated sodium hydrogen carbonate solution (3x) and dried (Na₂SO₄). The residue was suspended in 350 ml boiling CH₂Cl₂, cooled to 3 °C and filtered to yield 15.8 g (83%) of 3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ol, MS: 295 (M).

1.4

In analogy to examples 1.1 to 1.3, 4-bromo-benzoyl chloride and 3-fluoroanisole were converted to yield 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-ol, MS: 305 (M, 1Br).

1.5

- 5 In analogy to examples 1.1 to 1.3, 4-chloro-benzoyl chloride and 3-fluoroanisole were converted to yield 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-ol, MS: 261 (M, 1Cl).

1.6

- In analogy to examples 1.1 to 1.3, 4-fluoro-benzoyl chloride and 3-fluoroanisole were converted to yield 3-(4-Fluoro-phenyl)-benzo[d]isothiazol-6-ol as white solide, MS: 245
10 (M).

1.7

In analogy to example 1.2 to 1.3, 2-fluoro-4-methoxyacetophenone gave 3-Methyl-benzo[d]isothiazol-6-ol, MS: 166 (MH⁺).

Example 2

15 2.1

- 2.0 g (6.55 mmol) 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-ol in 30 ml acetone were treated with 5.89 g (42.5 mmol, 6.5 eq) K₂CO₃ and 1.9 ml (16.33 mmol, 2.5 eq) 1,4-dibromobutane. The reaction mixture was stirred at 45 °C for 8h and at RT over night, filtered and evaporated. The excess of dibromide was removed in vacuo and the crude
20 product was purified by column chromatography on silica gel with a gradient EtOAc:hexane 1:9 to 1:4 to yield 2.0 g (71%) 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole as yellow semisolid, MS: 439 (M, 2Br).

2.2

- In analogy to example 2.1, 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-ol and 1,2-
25 dibromoethane were converted to yield 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole as off-white solid, 105°C, MS: 411 (M, 2Br).

2.3

- In analogy to example 2.1, 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-ol and 1,3-dibromopropane were converted to yield 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-
30 benzo[d]isothiazole as yellow oil, MS: 425 (M, 2Br).

2.4

In analogy to example 2.1, 3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ol and 1,4-

dibromobutane were converted to yield 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole as light yellow solid, MS: 429 (M, 1Br).

2.5

In analogy to example 2.1, 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-ol and 1,4-
5 dibromobutane were converted to yield 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole as white solid, mp: 68-69°C, MS: 395 (M, 1Br, 1 Cl).

2.6

In analogy to example 2.1, 3-(4-fluoro-phenyl)-benzo[d]isothiazol-6-ol and 1,4-
dibromobutane were converted to yield 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-
10 benzo[d]isothiazole, MS: 379 (M, 1Br).

2.7

In analogy to example 2.1, 3-Methyl-benzo[d]isothiazol-6-ol and 1,3-dibromobutane were converted to yield 6-(4-Bromo-butoxy)-3-methyl-benzo[d]isothiazole as brown oil, MS: 300 (MH⁺, 1 Br).

15 Example 3

3.1

75 mg (0.18 mmol) 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole in 2.5 ml DMA were treated with 98 µl (0.54 mmol, 3eq) 5.6 M N,N-dimethylamine in ethanol for 35 h at RT. The solution was diluted with ether and a saturated aqueous
20 solution of NaHCO₃. The inorganic phase was extracted with ether, the combined organic phases were washed with brine and dried over Na₂SO₄. Column chromatography on silica gel with CH₂Cl₂:MeOH 8:1 yielded 63 mg (92%) [2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-dimethyl-amine as colorless oil, MS: 377(MH⁺, 1Br).

3.2

25 In analogy to example 3.1, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and N-Allylmethylamine were converted to yield Allyl-[2-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-methyl-amine as colorless oil, MS: 403(MH⁺, 1Br).

3.3

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-
30 benzo[d]isothiazole and N-Allylmethylamine were converted to yield Allyl-[3-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-methyl-amine as colorless oil, MS: 417(MH⁺, 1Br).

3.4

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and Dimethylamine were converted to yield [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-dimethyl-amine as white semisolid, MS: 391(MH⁺, 1Br).

3.5

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and N-(2-methoxyethyl)ethylamine were converted to yield (without extraction) [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-ethyl-(2-methoxy-ethyl)-amine hydrobromide as colorless oil, MS: 449(MH⁺, 1Br).

3.6

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and N-Allylmethylamine were converted to yield Allyl-[4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-methyl-amine as white semisolid, MS: 431(MH⁺, 1Br).

15 3.7

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and Dimethylamine were converted to yield [4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-dimethyl-amine as white semisolid, MS: 405(MH⁺, 1Br).

3.8

20 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and N-(2-Methoxyethyl)ethylamine were converted to {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine which was converted to the hydrochloride by treatment with HCl in methanol to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine hydrochloride as light
25 yellow oil, MS: 463(MH⁺, 1Br).

3.9

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and piperidine were converted to yield 3-(4-Bromo-phenyl)-6-(4-piperidin-1-yl-butoxy)-benzo[d]isothiazole as white solid, MS: 445(MH⁺, 1Br).

30 3.10

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 1-Methylpiperazine were converted to yield 3-(4-Bromo-phenyl)-6-[4-(4-methyl-piperazin-1-yl)-butoxy]-benzo[d]isothiazole as white solid, MS: 460(MH⁺, 1Br).

3.11

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 1-Acetylpiperazine were converted to yield 1-(4-(4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl)-piperazin-1-yl)-ethanone as white solid, MS: 488(MH⁺, 1Br).

3.12

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and Hydroxyethylamine were converted to yield 2-{3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propylamino}-ethanol as white semisolid, MS: 407 (MH⁺, 1Br).

3.13

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and Methyl-prop-2-ynyl-amine were converted to yield {3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-methyl-prop-2-ynyl-amine as colorless oil, MS: 415 (MH⁺, 1Br).

3.14

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and Prop-2-ynylamine were converted to yield {3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-prop-2-ynyl-amine as off-white semisolid, MS: 401 (MH⁺, 1Br).

3.15

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and Piperidine were converted to yield 3-(4-Bromo-phenyl)-6-(3-piperidin-1-yl-propoxy)-benzo[d]isothiazole as pink oil, MS: 431 (MH⁺, 1Br).

3.16

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and 1-Methyl-piperazine were converted to yield 3-(4-Bromo-phenyl)-6-[3-(4-methyl-piperazin-1-yl)-propoxy]-benzo[d]isothiazole as colorless oil, MS: 446 (MH⁺, 1Br).

3.17

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and 1-Piperazin-1-yl-ethanone were converted to yield 1-(4-{3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-piperazin-1-yl)-ethanone as colorless oil, MS: 474 (MH⁺, 1Br).

3.18

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and Ethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amine as white solid, MS: 405 (MH⁺, 1Br).

5 3.19

In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and Ethylamine were converted to yield {3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl}-ethyl-amine as white solid, MS: 391 (MH⁺, 1Br).

3.20

- 10 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[d]isothiazole and methylaminoethanol were converted to yield 2-({4-[3-(4-Fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol which was transferred into its salt by treatment with formic acid, MS: 375 (MH⁺).

3.21

- 15 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[d]isothiazole and N-(2-Methoxyethyl)methylamine were converted to yield {4-[3-(4-Fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine which was transferred into its salt by treatment with formic acid, MS: 389 (MH⁺).

3.22

- 20 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[d]isothiazole and N-(2-Methoxyethyl)ethylamine were converted to yield Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-amine which was transferred into its salt by treatment with formic acid, MS: 403 (MH⁺).

3.23

- 25 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[d]isothiazole and ethylaminoethanol were converted to yield 2-(Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol which was transferred into its salt by treatment with formic acid, MS: 389 (MH⁺).

3.24

- 30 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-methyl-benzo[d]isothiazole Ethylaminoethanol were converted to yield 2-{Ethyl-[4-(3-methyl-benzo[d]isothiazol-6-yloxy)-butyl]-amino}-ethanol as yellow oil, MS: 309 (MH⁺).

3.25

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-methyl-benzo[d]isothiazole N-(2-

Methoxyethyl)ethylamine were converted to yield Ethyl-(2-methoxy-ethyl)-[4-(3-methyl-benzo[d]isothiazol-6-yloxy)-butyl]-amine as yellow oil, MS: 323 (MH^+).

Example 4

1 eq of the corresponding bromide was treated with 3 eq of the corresponding amine in
5 DMF (1 ml/mmol bromide) in the presence of 1 eq diisopropylethylamine and a catalytic amount of NaI at RT until no starting material could be detected with HPLC. Formic acid was added and the crude materials were purified by preparative HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation the product was obtained as a mixture of amino-hydrobromides and formates.

10 4.1

According to the method in example 4, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 2-(methylamino)ethanol were converted to yield 2-[[2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-methyl-amino]-ethanol, MS: 408 (MH^+ , 1Br).

15 4.2

According to the method in example 4, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and N-(2-methoxyethyl)methylamine were converted to yield [2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-(2-methoxy-ethyl)-methyl-amine, MS: 422 (MH^+ , 1Br).

20 4.3

According to the method in example 4, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and morpholine were converted to yield 3-(4-Bromo-phenyl)-6-(2-morpholin-4-yl-ethoxy)-benzo[d]isothiazole, MS: 420 (MH^+ , 1Br).

4.4

25 According to the method in example 4, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 2-(ethylamino)ethanol were converted to yield 2-[[2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-ethyl]-ethyl-amino]-ethanol, MS: 422 (MH^+ , 1Br).

4.5

30 According to the method in example 4, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and pyrrolidine were converted to yield 3-(4-Bromo-phenyl)-6-(2-pyrrolidin-1-yl-ethoxy)-benzo[d]isothiazole, MS: 404 (MH^+ , 1Br).

4.6

According to the method in example 4, 6-(2-Bromo-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and azetidine were converted to yield 6-(2-Azetidin-1-yl-ethoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole, MS: 390 (MH⁺, 1Br).

5 4.7

According to the method in example 4, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and 2-(methylamino)ethanol were converted to yield 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-methyl-amino]-ethanol, MS: 422 (MH⁺, 1Br).

10 4.8

According to the method in example 4, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and N-(2-methoxyethyl)methylamine were converted to yield [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-(2-methoxy-ethyl)-methylamine, MS: 436 (MH⁺, 1Br).

15 4.9

According to the method in example 4, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and morpholine were converted to yield 3-(4-Bromo-phenyl)-6-(3-morpholin-4-yl-propoxy)-benzo[d]isothiazole, MS: 434 (MH⁺, 1Br).

4.10

20 According to the method in example 4, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and 2-(ethylamino)ethanol were converted to yield 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-ethyl-amino]-ethanol, MS: 436 (MH⁺, 1Br).

4.11

25 According to the method in example 4, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and pyrrolidine were converted to yield 3-(4-Bromo-phenyl)-6-(3-pyrrolidin-1-yl-propoxy)-benzo[d]isothiazole, MS: 418 (MH⁺, 1Br).

4.12

30 According to the method in example 4, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and azetidine were converted to yield 6-(3-Azetidin-1-yl-propoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole, MS: 404 (MH⁺, 1Br).

4.13

According to the method in example 4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 2-(methylamino)ethanol were converted to yield 2-[[4-[3-(4-

Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-methyl-amino]-ethanol, MS: 436 (MH⁺, 1Br).

4.14

According to the method in example 4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and N-(2-methoxyethyl)methylamine were converted to yield [4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-(2-methoxy-ethyl)-methyl-amine, MS: 450 (MH⁺, 1Br).

4.15

According to the method in example 4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and morpholine were converted to yield 3-(4-Bromo-phenyl)-6-(4-morpholin-4-yl-butoxy)-benzo[d]isothiazole, MS: 448 (MH⁺, 1Br).

4.16

According to the method in example 4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 2-(ethylamino)ethanol were converted to yield 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-ethyl-amino]-ethanol, MS: 450 (MH⁺, 1Br).

4.17

According to the method in example 4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and pyrrolidine were converted to yield 3-(4-Bromo-phenyl)-6-(4-pyrrolidin-1-yl-butoxy)-benzo[d]isothiazole, MS: 432 (MH⁺, 1Br).

4.18

According to the method in example 4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and azetidine were converted to yield 6-(4-Azetidin-1-yl-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole, MS: 418 (MH⁺, 1Br).

25 Example 5

1 eq of the corresponding bromide was treated with 3 eq of the corresponding amine in DMA (1 ml/0.25 mmol bromide) at RT until no starting material could be detected with HPLC. Formic acid was added and the crude materials were purified by preparative HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation the product was obtained as a mixture of amino-hydrobromides and formates.

5.1

According to the method in example 5, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-

benzo[d]isothiazole and diethanolamine were converted to yield 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-(2-hydroxy-ethyl)-amino]-ethanol as white semisolid, MS: 451 (MH⁺, 1Br).

5.2

- 5 According to the method in example 5, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole and bis(2-methoxyethyl)amine were converted to yield [3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-bis-(2-methoxy-ethyl)-amine as orange oil, MS: 479 (MH⁺, 1Br).

5.2

- 10 According to the method in example 5, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and diethanolamine were converted to yield 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol as white semisolid, MS: 465(MH⁺, 1Br).

5.3

- 15 According to the method in example 5, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and bis(2-methoxyethyl)amine were converted to yield [4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-bis-(2-methoxy-ethyl)-amine as light yellow oil, MS: 493 (MH⁺, 1Br).

Example 6

- 20 1 eq of the corresponding bromide was treated with 3 eq of the corresponding amine in DMA (1 ml/0.25mmol bromide) at RT until no starting material could be detected with HPLC. Formic acid was added and the crude materials were purified by prep HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After separation, the product was extracted with EtOAc and sat. NaHCO₃/H₂O to isolate the
25 product as free amine. The hydrochloride salt was optionally obtained after addition of 1N HCl/MeOH and evaporation of the solvent.

6.1

- According to the method in example 6, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and 1,1-dimethylpropargylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(1,1-dimethyl-prop-2-ynyl)-amine as
30 colorless oil, MS: 443(MH⁺, 1Br).

6.2

According to the method in example 6, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-

benzo[d]isothiazole and ethanolamine were converted to yield 2-{4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butylamino}-ethanol as white solid, MS: 421(MH⁺, 1Br).

6.3

According to the method in example 6, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and Methyl-prop-2-ynyl-amine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-prop-2-ynyl-amine as light yellow oil, MS: 429(MH⁺, 1Br).

6.4

According to the method in example 6, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole and propargylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-prop-2-ynyl-amine hydrochloride as off-white solid, MS: 415(MH⁺, 1Br).

Example 7

A solution of 0.25 mmol (1 equivalent) of the corresponding bromide in 0.7 ml dry DMA was treated with a solution of 0.5 mmol (2 equivalents) of the corresponding secondary amine in 0.15 ml dry DMA at room temperature. After 16 h, 2 equivalents of secondary amine were added again to the solution. The reaction mixture was allowed to stand over night at room temperature, treated with 0.2 ml formic acid and purified by preparative HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation, the product was obtained as a mixture of amino hydrobromide and formate. Optionally the product was extracted with EtOAc and sat. NaHCO₃/H₂O to obtain the corresponding free amine.

7.1

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Diethylamine were converted to yield Diethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine, MS: 423 (MH⁺).

7.2

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Piperidine were converted to yield 6-(4-Piperidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole, MS: 435 (MH⁺).

7.3

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Allylmethylamine were converted to yield Allyl-methyl-

{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine, MS: 421 (MH⁺).

7.4

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Trimethylene imine were converted to yield 6-(4-Azetidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole, MS: 407 (MH⁺).

7.5

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Ethylamino ethanol were converted to yield 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol, MS: 439 (MH⁺).

7.6

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Diethanoleamine were converted to yield 2-((2-Hydroxy-ethyl)-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol, MS: 455 (MH⁺).

7.7

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole and Dimethylamine were converted to yield Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine, MS: 395 (MH⁺).

7.8

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and N-Allylmethylamine were converted to yield Allyl-{4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amine, MS: 387 (MH⁺, 1Cl).

25 7.9

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and 2-Ethylamino-ethanol were converted to yield 2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol, MS: 405 (MH⁺, 1Cl).

30 7.10

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and Azetidine were converted to yield 6-(4-Azetidin-1-yl-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole, MS: 373 (MH⁺, 1Cl).

7.11

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and Piperidine were converted to yield 3-(4-Chloro-phenyl)-6-(4-piperidin-1-yl-butoxy)-benzo[d]isothiazole, MS: 401 (MH^+ , 1Cl).

5 7.12

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and N-(2-Methoxyethyl)ethylamine were converted to yield {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine, MS: 419 (MH^+ , 1Cl).

10 7.13

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and Diethylamine were converted to yield {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine, MS: 389 (MH^+ , 1Cl).

7.14

15 According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and N-Methylcyclopropanemethylamine were converted to yield {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-cyclopropylmethyl-methyl-amine, MS: 401 (MH^+ , 1Cl).

7.15

20 According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and N-Isopropylmethylamine were converted to yield {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-isopropyl-methyl-amine, MS: 389 (MH^+ , 1Cl).

7.16

25 According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and Pyrrolidine were converted to yield 3-(4-Chloro-phenyl)-6-(4-pyrrolidin-1-yl-butoxy)-benzo[d]isothiazole, MS: 387 (MH^+ , 1Cl).

7.17

30 According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and N-(2-Methoxyethyl)Methylamine were converted to yield {4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine, MS: 405 (MH^+ , 1Cl).

7.18

According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-

benzo[d]isothiazole and 2-(Methylamino)Ethanol were converted to yield 2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol, MS: 391 (MH⁺, 1Cl).

7.19

- 5 According to the method in example 7, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole and (R)-3-Hydroxypyrrolidine were converted to yield (3R)-1-{4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-pyrrolidin-3-ol, MS: 403 (MH⁺, 1Cl).

Example 8

10 8.1

To 4.4 g (10 mmol) 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole in 400 ml CH₂Cl₂, 24 g potassium permanganate on silica gel were added, which were prepared from 7 g KMnO₄, 20g silica gel and 10 ml water prior to use. The mixture was stirred at 50°C for 2 days, filtered and the crude material purified by column

- 15 chromatography with EtOAc:hexane 1:4 to give 1.6 g (34%) 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide as off-white solid, MS: 473 (MH⁺, 1Br).

8.2

In analogy to example 8.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole was converted to 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-

- 20 benzo[d]isothiazole 1,1-dioxide as off-white solid, MS: 457 (M, 2Br).

8.3

In analogy to example 8.1, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole was converted to 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide as light brown solid, which was used in subsequent

- 25 reaction steps without further purification.

8.4

In analogy to example 8.1, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole was converted to 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide as light yellow solid, MS: 426 (M, 1Br, 1 Cl).

30 Example 9

1 eq of the corresponding bromide was treated with 3 eq of the corresponding amine in DMA (1 ml/0.25 mmol bromide) at RT until no starting material could be detected with HPLC. Formic acid was added and the crude materials were purified by preparative HPLC

[RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation the product was obtained as a mixture of amino-hydrobromides and formates.

9.1

- 5 According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and dimethylamine were converted to yield 3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-dimethyl-amine as yellow solid, MS: 423 (MH⁺, 1Br).

9.2

- 10 According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and N-allylmethylamine were converted to yield Allyl-(3-[3-(4-bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-methyl-amine as yellow solid, MS: 449 (MH⁺, 1Br).

9.3

- 15 According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and azetidine were converted to yield 6-(3-Azetidin-1-yl-propoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide as yellow solid, MS: 435 (MH⁺, 1Br).

9.4

- 20 According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and N-Methylproylamine were converted to yield 3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-methyl-propyl-amine as yellow solid, MS: 451 (MH⁺, 1Br).

9.5

- 25 According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and 2-Ethylaminoethanol were converted to yield 2-((3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-ethyl-amino)-ethanol as yellow solid, MS: 467(MH⁺, 1Br).

9.6

- 30 According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and diethylamine were converted to yield 3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-propyl)-diethyl-amine as yellow solid, MS: 451 (MH⁺, 1Br).

9.7

According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and N-(2-methoxyethyl)methylamine were converted to yield {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-propyl}-(2-methoxy-ethyl)-methyl-amine as yellow solid, MS: 467 (MH⁺, 1Br).

9.8

According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and 2-methylaminoethanol were converted to yield 2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-propyl}-methyl-amino)-ethanol as yellow solid, MS: 453 (MH⁺, 1Br).

9.9

According to the method in example 9, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[d]isothiazole 1,1-dioxide and N-(2-methoxyethyl)ethylamine were converted to yield {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-propyl}-ethyl-(2-methoxy-ethyl)-amine as yellow solid, MS: 481 (MH⁺, 1Br).

9.10

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and dimethylamine were converted to yield (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-dimethyl-amine as yellow solid, MS: 437 (MH⁺, 1Br).

9.11

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and N-allylmethylamine were converted to yield Allyl-(4-[3-(4-bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-methyl-amine as yellow solid, MS: 463 (MH⁺, 1Br).

9.12

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and azetidine were converted to yield 6-(4-Azetidin-1-yl-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide as yellow solid, MS: 449 (MH⁺, 1Br).

9.13

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and N-Methylproylamine were converted to yield {4-[3-

(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-methyl-propyl-amine as yellow solid, MS: 465 (MH⁺, 1Br).

9.14

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and 2-Ethylaminoethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol as yellow solid, MS: 481 (MH⁺, 1Br).

9.15

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and diethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine as yellow solid, MS: 465 (MH⁺, 1Br).

9.16

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and N-(2-methoxyethyl)methylamine were converted to yield {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine as yellow solid, MS: 481 (MH⁺, 1Br).

9.17

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and 2-methylaminoethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amino)-ethanol as yellow solid, MS: 467 (MH⁺, 1Br).

9.18

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and N-(2-methoxyethyl)ethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine as yellow solid, MS: 495 (MH⁺, 1Br).

9.19

According to the method in example 9, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[d]isothiazole 1,1-dioxide and bis(2-methoxyethyl)amine were converted to yield (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-bis-(2-methoxy-ethyl)-amine, which was converted with HCl/MeOH to yield the corresponding hydrochloride as yellow semisolid, MS: 525(MH⁺, 1Br).

Example 10

A solution of 0.25 mmol (1 equivalent) of the corresponding bromide in 0.7 ml dry DMA was treated with a solution of 0.5 mmol (2 equivalents) of the corresponding secondary amine in 0.15 ml dry DMA at room temperature. After 16 h, additional 2 equivalents of
5 secondary amine were added to the solution. The reaction mixture was allowed to stand over night at room temperature, treated with 0.2 ml formic acid and purified by preparative HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation, the product was obtained as a mixture of amino hydrobromide and formate.

10 10.1

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and Allylmethylamine were converted to yield Allyl-{4-[1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-amine, MS: 453 (MH⁺).

15 10.2

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and 2-Ethylaminoethanol were converted to yield 2-({4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol, MS: 471 (MH⁺).

20 10.3

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and Diethylamine were converted to yield {4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine, MS: 455 (MH⁺).

25 10.4

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and Dimethylamine were converted to yield {4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-dimethyl-amine, MS: 427 (MH⁺).

30 10.5

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and Piperidine were converted to yield 6-(4-Piperidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide, MS: 467 (MH⁺).

10.6

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and Trimethyleneamine were converted to yield 6-(4-Azetidin-1-yl-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide,
5 MS: 439 (MH⁺).

10.7

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole 1,1-dioxide and Diethanolamine were converted to yield 2-
[4-[1,1-Dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl]-
10 (2-hydroxy-ethyl)-amino]-ethanol, MS: 487 (MH⁺).

10.8

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Piperidine were converted to yield 3-(4-Chloro-phenyl)-6-(4-piperidin-1-yl-butoxy)-benzo[d]isothiazole 1,1-dioxide, MS: 433 (MH⁺,
15 1Cl).

10.9

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Allyl-methyl-amine were converted to yield Allyl-{4-[3-(4-chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-
20 amine, MS: 419 (MH⁺, 1Cl).

10.10

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Azetidine were converted to yield 6-(4-Azetidin-1-yl-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide, MS: 405 (MH⁺, 1Cl).

25 10.11

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Ethyl-(2-methoxy-ethyl)-amine were converted to yield {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine, MS: 451 (MH⁺, 1Cl).

30 10.12

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and 2-Ethylamino-ethanol were converted to yield 2-({4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol, MS: 437 (MH⁺, 1Cl).

10.13

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Diethyl-amine were converted to yield {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-diethyl-amine,
5 MS: 421 (MH⁺, 1Cl).

10.14

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and 2-Methylamino-ethanol were converted to yield 2-({4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-
10 amino)-ethanol, MS: 423 (MH⁺, 1Cl).

10.15

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and (2-Methoxy-ethyl)-methyl-amine were converted to yield {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-
15 methoxy-ethyl)-methyl-amine, MS: 437 (MH⁺, 1Cl).

10.16

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Pyrrolidine were converted to yield 3-(4-Chloro-phenyl)-6-(4-pyrrolidin-1-yl-butoxy)-benzo[d]isothiazole 1,1-dioxide, MS: 419 (MH⁺,
20 1Cl).

10.17

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and (3R)-Pyrrolidin-3-ol were converted to yield 1-{4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-pyrrolidin-3-ol,
25 MS: 435 (MH⁺, 1Cl).

10.18

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Isopropyl-methyl-amine were converted to yield {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-1l 6-benzo[d]isothiazol-6-yloxy]-butyl}-isopropyl-
30 methyl-amine, MS: 421 (MH⁺, 1Cl).

10.19

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and Cyclopropylmethyl-methyl-amine were converted to

yield {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-cyclopropylmethyl-methyl-amine, MS: 433 (MH^+ , 1Cl).

10.20

According to the method in example 10, 6-(4-Bromo-butoxy)-3-(4-chloro-phenyl)-benzo[d]isothiazole 1,1-dioxide and N-Methylpropylamine were converted to yield {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-propyl-amine, MS: 421 (MH^+ , 1Cl).

Example 11

11.1

To a suspension of 36 mg (0.38 mmol, 2.2 eq) Hydrogen peroxide-Urea adduct in 0.2 ml CH_2Cl_2 , 28 mg (0.19 mmol, 1.1 eq) phthalic anhydride were added and stirred for 15 min at RT. 75 mg (0.17 mmol, 1.0 eq) {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-dimethyl-amine in 0.3 ml CH_2Cl_2 were added and the mixture was stirred at RT for 2h. 1M aqueous K_2CO_3 solution was added (pH 7) and the inorganic phase was extracted with CH_2Cl_2 . The organic phases were washed with water and brine and dried over Na_2SO_4 . Column chromatography on silica gel with CH_2Cl_2 :MeOH 8:1 yielded 48 mg (62%) {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-dimethyl-amine N-oxide as light yellow semisolid, MS: 453(MH^+ , 1Br).

11.2

In analogy to example 11.1, {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine was converted to yield {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine N-oxide as light yellow semisolid, MS: 497 (MH^+ , 1Br).

11.3

In analogy to example 11.1, (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-bis-(2-methoxy-ethyl)-amine was converted to yield (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-bis-(2-methoxy-ethyl)-amine N-oxide as yellow semisolid, MS: 541 (MH^+ , 1Br).

11.4

In analogy to example 11.1, {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine was converted to {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine N-oxide which was dissolved in THF and treated with 1M Bu_4NF in THF over

night (see example 36.1) to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol N-oxide as off-white semisolid, MS: 465 (MH^+ , 1Br).

Example 12

12.1

5 5.0 g (16.33 mmol) 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-ol in 8 ml pyridine were treated with 3 ml (17.96 mmol) trifluoromethane sulfonic anhydride at 0°C. The solution was stirred for 1.5 h at RT. The solution was diluted with ether and water, the phases were separated and the inorganic phase was extracted with ether. The combined organic phases were washed with 2M HCl and brine and dried over Na_2SO_4 . Evaporation yielded 7.21
10 crude Trifluoro-methanesulfonic acid 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl ester as yellow oil, MS: 437 (M, 1Br).

12.2

In analogy to example 12.1, 3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ol and trifluoromethane sulfonic anhydride were converted to yield Trifluoro-methanesulfonic
15 acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester as light brown solid, MS: 427 (M).

12.3

In analogy to example 12.1, 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-ol and trifluoromethane sulfonic anhydride were converted to yield Trifluoro-methanesulfonic
20 acid 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl ester as orange oil, MS: 393 (MH^+ , 1Cl).

Example 13

13.1

3.0 g (6.846 mmol) Trifluoro-methanesulfonic acid 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl ester were suspended three times with toluene prior to the
25 reaction, then dissolved in 25 ml THF and added to 240 mg (0.34 mmol) $PdCl_2(PPh_3)_2$. To this suspension 0.67 ml (7.19 mmol) 4-pentyn-1-ol in 1 ml THF were added, followed by 54 mg (0.205 mmol) triphenylphosphine in 1 ml THF and 2.9 ml (20.54 mmol) triethylamine. The solution was stirred for 20 min at RT prior to the addition of 13 mg (0.07 mmol) CuI. The solution was heated to 50°C overnight. Further 132 μ l (1.4 mmol) 4-
30 pentyn-1-ol in 0.8 ml THF were added in two portions and stirring at 50 °C was continued for 5 h and at RT over night. The reaction mixture was diluted with EtOAc and 1M $KHSO_4$. The inorganic phase was extracted with EtOAc and the combined organic phases were washed with water and brine and dried over Na_2SO_4 . Column chromatography with a

gradient hexane:EtOAc 2:1 to 1:1 gave 1.83 g (72%) 5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-yn-1-ol as light yellow oil, MS: 371 (M, 1Br).

13.2

In analogy to example 13.1, Trifluoro-methanesulfonic acid 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl ester and 2-propyn-1-ol were converted to yield 3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol as light brown solid, MS: 343 (MH⁺, 1Br).

13.3

In analogy to example 13.1, Trifluoro-methanesulfonic acid 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl ester and 3-butyne-1-ol were converted to yield 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol as orange solid, MS: 357 (M, 1Br).

13.4

In analogy to example 13.1, Trifluoro-methanesulfonic acid 3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl ester and 2-Methyl-3-butyne-2-ol were converted to yield 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-2-methyl-but-3-yn-2-ol as light yellow semisolid, MS: 371 (M, 1Br).

Example 14

14.1

To 3.3 g (7.72 mmol) Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester in 20 ml piperidine 447.0 mg (3.9 mmol) tetrakis(triphenylphosphine)palladium and 73.6 mg (3.9 mmol) CuI were added. At 80°C 1.43 ml (15.45 mmol) 4-pentyne-1-ol were added slowly, and the solution was stirred for an additional hour. The cooled solution was added to ice water, acidified with 2M HCl and extracted with ether. The combined organic phases were washed with water and dried over Na₂SO₄. Column chromatography on silica gel with CH₂Cl₂:MeOH 30:1 yielded 2.7 g (96%) 5-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-yn-1-ol as orange oil, MS: 361(M).

14.2

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and 2-propyn-1-ol were converted to yield 3-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol as off-white solid, MS: 333(M).

14.3

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-

benzo[d]isothiazol-6-yl ester and 3-butyne-1-ol were converted to yield 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol as orange semisolid, MS: 347 (M).

14.4

- 5 In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl ester and 2-propyne-1-ol were converted to yield 3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol as yellow solid, MS: 299 (M, 1 Cl).

14.5

- 10 In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl ester and 3-butyne-1-ol were converted to yield 4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol as yellow solid, mp: 128-129 °C, MS: 313 (M, 1 Cl).

14.6

- 15 In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl ester and 4-pentyne-1-ol were converted to yield 5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-yn-1-ol as yellow oil, MS: 327 (M, 1 Cl).

Example 15

15.1

- 20 365 mg (1.02 mmol) 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol in 7 ml CH₂Cl₂ were treated with 94 µl (1.21 mmol) methane sulfonic acid chloride, 0.13 ml (1.53 mmol) pyridine and 135 mg (1.2 mmol) DMAP at 0 °C. The solution was stirred at RT over night, was diluted with water and acidified with 10% aq. KHSO₄. The inorganic phase was extracted with CH₂Cl₂. The combined organic phases were washed with NaHCO₃ solution, brine and dried over Na₂SO₄. Evaporation gave 446 mg crude
- 25 Methanesulfonic acid 4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl ester, MS: 436 (MH⁺).

15.2

- 30 In analogy to example 15.1, 3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 3-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester, which was directly subjected to the following reaction.

15.3

In analogy to example 15.1, 5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 5-[3-(4-

bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester, MS: 450 (MH⁺, 1Br).

15.4

In analogy to example 15.1, 3-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester, which
5 was directly subjected to the following reaction.

15.5

In analogy to example 15.1, 5-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester light
10 brown oil, MS: 369(M).

15.6

In analogy to example 15.1, 3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester as a brown oil, MS: 377 (M,
15 1Cl).

15.7

In analogy to example 15.1, 4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl ester as a light yellow oil, MS: 391 (M,
20 1Cl).

15.8

In analogy to example 15.1, 5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-yn-1-ol and methane sulfonyl chloride were converted to yield Methanesulfonic acid 5-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester as a light yellow oil, MS: 405
25 (M, 1Cl).

Example 16

16.1

73 mg (0.16 mmol) Methanesulfonic acid 4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl ester in 1 ml DMA were treated with 35 μ l (0.5 mmol) azetidine for 48 h.
30 The solution was diluted with ether and aqueous Na₂CO₃, the phases were separated and the inorganic layer was extracted with ether. The combined organic phases were washed with brine, dried over Na₂SO₄. Column chromatography on silica gel with a gradient

CH₂Cl₂ to CH₂Cl₂:MeOH 8:1 gave 22.2 mg (33%) 6-(4-Azetidin-1-yl-but-1-ynyl)-3-(4-bromo-phenyl)-benzo[d]isothiazole as brown oil, MS: 397(MH⁺, 1Br).

16.2

In analogy to example 16.1, Methanesulfonic acid 4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl ester and 2-(Methylamino)ethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-methyl-amino)-ethanol as brown oil, MS: 415(MH⁺, 1Br).

16.3

In analogy to example 16.1, Methanesulfonic acid 5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and Allylmethylamine were converted to yield Allyl-{5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-methyl-amine as light yellow solid, MS: 425 (MH⁺, 1Br).

16.4

In analogy to example 16.1, Methanesulfonic acid 5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and 2-Methylaminoethanol were converted to yield 2-({5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-methyl-amino)-ethanol as off-white semisolid, MS: 429(MH⁺, 1Br).

16.5

In analogy to example 16.1, Methanesulfonic acid 5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and 2-Ethylaminoethanol were converted to yield 2-({5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol as off-white semisolid, MS: 443 (MH⁺, 1Br).

16.6

In analogy to example 16.1, Methanesulfonic acid 5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and N-(2-Methoxyethyl)methylamine were converted to yield {5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-(2-methoxy-ethyl)-methyl-amine as off-white semisolid, MS: 443 (MH⁺, 1Br).

16.7

In analogy to example 16.1, Methanesulfonic acid 5-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and Dimethylamine were converted to yield {5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-dimethyl-amine as off-white semisolid, MS: 399 (MH⁺, 1Br).

16.8

In analogy to example 16.1, Methanesulfonic acid 5-[3-(4-bromo-phenyl)-

benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and azetidine were converted to yield 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-bromo-phenyl)-benzo[d]isothiazole as orange semisolid, MS: 411 (MH^+ , 1Br).

16.9

- 5 In analogy to example 16.1, Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and Allylmethylamine (2x 2 equivalents) were converted to yield Allyl-{3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-methyl-amine as brown oil, MS: 353 (MH^+ , 1Cl).

16.10

- 10 In analogy to example 16.1, Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and 2-Ethylaminoethanol (2x 2 equivalents) were converted to yield 2-({3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-ethyl-amino)-ethanol, MS: 371 (MH^+ , 1Cl).

16.11

- 15 In analogy to example 16.1, Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and Diethanolamine (2x 2 equivalents) were converted to yield 2-({3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-(2-hydroxy-ethyl)-amino)-ethanol as white solid, MS: 387 (MH^+ , 1Cl).

16.12

- 20 In analogy to example 16.1, Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and Dimethylamine (10 equivalents of a 5.6 M EtOH solution) were converted to yield {3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-dimethyl-amine as light brown solid, MS: 327 (MH^+ , 1Cl).

Example 17

25 17.1

- 150 mg (0.34 mmol) Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester in 1.5 ml DMF were treated with 140 μ l (1.7 mmol) N-allylmethylamine at 50°C for 4h. The solution was diluted with ether and water, the organic phase was washed with 0.5 M NaOH, and dried over Na_2SO_4 . Column chromatography on silica gel gave 80 mg (56%) 2-(Methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amino)-ethanol as light yellow, MS: 419(MH^+).

17.2

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-bromo-phenyl)-

benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and 2-methylaminoethanol were converted to yield 2-({3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-methyl-amino)-ethanol as light brown semisolid, MS: 401(MH⁺, 1Br).

17.3

- 5 In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and 2-ethylaminoethanol were converted to yield 2-({3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-ethyl-amino)-ethanol as light brown semisolid, MS: 415(MH⁺, 1Br).

17.4

- 10 In analogy to example 17.1, Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and N-allylmethylamine were converted to yield Allyl-methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine as light yellow oil, MS: 415(MH⁺).

17.5

- 15 In analogy to example 17.1, Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and 2-Methoxyethylmethylamine were converted to yield (2-Methoxy-ethyl)-methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine as orange oil, MS: 433 (MH⁺).

17.6

- 20 In analogy to example 17.1, Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and dimethylamine were converted to yield Dimethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine as off-white solid, MS: 389 (MH⁺).

17.7

- 25 In analogy to example 17.1, Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and 2-Ethylaminoethanol were converted to yield 2-(Ethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amino)-ethanol as light yellow oil, MS: 433 (MH⁺).

17.8

- 30 In analogy to example 17.1, Methanesulfonic acid 5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and azetidine were converted to yield 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole as orange oil, MS: 401 (MH⁺).

17.9

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and 2-Methylaminoethanol were converted to yield 2-(Methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol as light brown oil, MS: 391 (MH⁺).

17.10

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and 2-ethylaminoethanol were converted to yield 2-(Ethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol as light brown oil, MS: 405 (MH⁺).

17.11

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and N-allylmethylamine were converted to yield Allyl-methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine as orange oil, MS: 387 (MH⁺).

17.12

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and azetidine were converted to yield 6-(3-Azetidin-1-yl-prop-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole as light brown oil, MS: 373 (MH⁺).

17.13

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and 2-Methoxyethylmethylamine were converted to yield (2-Methoxy-ethyl)-methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine as light brown oil, MS: 405 (MH⁺).

17.14

In analogy to example 17.1, Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl ester and Dimethylamine were converted to yield Dimethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine as light brown oil, MS: 361 (MH⁺).

Example 18

A solution of 0.25 mmol (1 equivalent) of the corresponding methanesulfonic acid in 0.7 ml dry DMA was treated with a solution of 0.5 mmol (2 equivalents) of the corresponding secondary amine in 0.15 ml dry DMA at room temperature. After 16 h, 2 equivalents of

secondary amine were added again to the solution. The reaction mixture was allowed to stand over night at room temperature, treated with 0.2 ml formic acid and purified by preparative HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation, the product was obtained as a mixture of amino
5 formate and methanesulfonate.

18.1

According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and N-Dimethylamine were converted to yield
10 {5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-dimethyl-amine, MS: 355 (MH⁺).

18.2

According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and Piperidine were converted to yield 3-(4-Chloro-phenyl)-6-(5-piperidin-1-yl-pent-1-ynyl)-benzo[d]isothiazole, MS: 395 (MH⁺).

15 18.3

According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and Azetidine were converted to yield 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-chloro-phenyl)-benzo[d]isothiazole, MS: 367 (MH⁺).

18.4

20 According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and N-Methylallylamine were converted to yield Allyl-{5-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-methyl-amine, MS: 381 (MH⁺).

18.5

25 According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and Diethylamine were converted to yield {5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-diethyl-amine, MS: 383 (MH⁺).

18.6

30 According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and Diethanolamine were converted to yield 2-[[5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl]-(2-hydroxy-ethyl)-amino]-ethanol, MS: 415 (MH⁺).

18.7

According to the method in example 18, Methanesulfonic acid 5- [3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl ester and 2-(Ethylamino)ethanol were converted to yield 2-({5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol, MS: 399 (MH⁺).

Example 19

19.1

100 mg (0.28 mmol)) 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol in 2 ml CH₂Cl₂ were treated with 53 µl (0.31 mmol) Huenig's base at 51 µl (0.31 mmol) trifluoromethane sulfonic acid anhydride at -15 °C. The solution was stirred at that temperature for 1h, additional 10 µl trifluoromethane sulfonic acid anhydride were added and stirring continued for 1h. 48 µl (0.28 mmol) Huenig's base and 75 µl (0.84 mmol) N-(2-Methoxyethyl)methylamine were added at -15°C. The solution was warmed to RT and stirred for 1.5 h. The solution was diluted with CH₂Cl₂ and washed with Na₂CO₃ solution, brine and dried with Na₂SO₄. Column chromatography gave 66.1 mg (56%) 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl)-(2-methoxy-ethyl)-methyl-amine as brown oil, MS: 429 (MH⁺, 1Br).

19.2

In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and Dimethylamine were converted to yield 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl)-dimethyl-amine as light brown semisolid, MS: 385 (MH⁺, 1Br).

19.3

In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and N-Allylmethylamine were converted to yield Allyl-4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl)-methyl-amine as brown oil, MS: 411 (MH⁺, 1Br).

19.4

In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and 2-(Ethylamino)ethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-ethyl-amino)-ethanol as brown oil, MS: 429 (MH⁺, 1Br).

19.5

In analogy to example 19.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and Dimethylamine were converted to yield Dimethyl-4-[3-(4-

trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amine as brown oil, MS: 375 (MH⁺).

19.6

In analogy to example 19.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and N-allylmethylamine were converted to yield Allyl-methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amine as yellow oil, MS: 401 (MH⁺).

19.7

In analogy to example 19.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and 2-Ethylaminoethanol were converted to yield 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amino)-ethanol as light yellow oil, MS: 419 (MH⁺).

19.8

In analogy to example 19.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and 2-Methylaminoethanol were converted to yield 2-(Methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amino)-ethanol as yellow oil, MS: 405 (MH⁺).

19.9

In analogy to example 19.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and 2-Methoxyethylmethylamine were converted to yield (2-Methoxyethyl)-methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amine as light brown oil, MS: 419 (MH⁺).

19.10

In analogy to example 19.1, 4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol and 2-Ethylaminoethanol were converted to yield 2-({4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-ethyl-amino)-ethanol as light yellow oil, MS: 358 (MH⁺, 1Cl).

Example 20

20.1

500 mg (1.5 mmol) 3-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol in 25 ml MeOH were hydrogenated in the presence of 120 mg PtO₂. H₂O, filtration over silica gel and evaporation yielded 360 mg (71%) 3-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propan-1-ol, which was directly subjected to the following reaction.

20.2

In analogy to example 20.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol was converted to yield 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol, MS: 351 (M).

5 20.3

In analogy to example 20.1, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl ester in EtOH/dioxane was converted to yield Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester as a dark brown oil, MS: 396 (MH⁺, 1Cl).

10 20.4

In analogy to example 20.1, 3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-yn-1-ol in EtOH/dioxane was converted to yield 3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-propan-1-ol as a brown oil, MS: 303 (M, 1Cl).

20.5

15 In analogy to example 20.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-but-3-yn-1-ol was converted to yield 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol as yellow oil, MS: 361 (M, 1Br).

Example 21

21.1

20 360 mg (1.07 mmol) 3-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propan-1-ol in 3 ml CH₂Cl₂ were treated with 100 μ l (1.29 mmol) methanesulfonic acid chloride and 450 μ l (3.23 mmol) triethylamine at 0°C. The solution was stirred at RT for 2h, diluted with CH₂Cl₂ and washed with 1N HCl and dried over Na₂SO₄. Evaporation gave 350 mg (79%) crude Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl ester, which was directly subjected to the following reaction.

21.2

In analogy to example 21.1, 4-[3-(4-Trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and methanesulfonic acid chloride were converted to yield Methanesulfonic acid 4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester, MS: 430 (MH⁺).

30

21.3

In analogy to example 21.1, 3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-propan-1-ol and methanesulfonic acid chloride were converted to Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-propyl ester as a orange oil, MS: 381 (M, 1Cl)

Example 22

22.1

In analogy to example 17.1; Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl ester and 2-ethylaminoethanol were converted to yield 2-
5 (Ethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-amino)-ethanol as light brown oil, MS: 409 (MH⁺).

22.2

In analogy to example 17.1; Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl ester and Dimethylamine were converted to yield
10 Dimethyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-amine as light brown oil, MS: 365 (MH⁺).

22.3

In analogy to example 17.1; Methanesulfonic acid 3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl ester and 2-Methylaminoethanol were converted to yield
15 2-(Methyl-{3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-amino)-ethanol as light brown oil, MS: 395 (MH⁺).

22.4

In analogy to example 17.1; Methanesulfonic acid 4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and 2-Ethylaminoethanol were converted to yield 2-
20 (Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-amino)-ethanol as light yellow oil, MS: 423 (MH⁺).

22.5

In analogy to example 17.1; Methanesulfonic acid 4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and Dimethylamine were converted to yield
25 Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-amine as yellow oil, MS: 379 (MH⁺).

22.6

In analogy to example 17.1; Methanesulfonic acid 4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and 2-Methylaminoethanol were converted to yield 2-
30 (Methyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-amino)-ethanol as light yellow oil, MS: 409 (MH⁺).

22.7

In analogy to example 17.1; Methanesulfonic acid 3-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-propyl ester and 2-Ethylaminoethanol were converted to yield 2-

((3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-propyl)-ethyl-amino)-ethanol as yellow oil, MS: 375 (MH^+ , 1Cl).

Example 23

A solution of 0.25 mmol (1 equivalent) of the corresponding methanesulfonic acid in 0.7 ml dry DMA was treated with a solution of 0.5 mmol (2 equivalents) of the corresponding secondary amine in 0.15 ml dry DMA at room temperature. After 16 h, 2 equivalents of secondary amine were added again to the solution. The reaction mixture was allowed to stand over night at room temperature, treated with 0.2 ml formic acid and purified by preparative HPLC [RP-18, acetonitrile (0.1 % $HCOOH$)/water (0.1 % $HCOOH$), 10 % to 95 % acetonitrile]. After evaporation, the product was obtained as a mixture of amino formate and methanesulfonate.

23.1

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and N-Methylallylamine were converted to yield Allyl-4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl-methyl-amine, MS: 371 (MH^+ , 1Cl).

23.2

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and Diethylamine were converted to yield 4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl-diethyl-amine, MS: 373 (MH^+ , 1Cl).

23.3

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and N-Methylpropylamine were converted to yield 4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl-methyl-propyl-amine, MS: 373 (MH^+ , 1Cl).

23.4

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and Diethanolamin were converted to yield 2-[[4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol, MS: 405 (MH^+ , 1Cl).

23.5

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and 2-Ethylaminoethanol were converted to yield 2-

((4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl)-ethyl-amino)-ethanol, MS: 389 (MH⁺, 1Cl).

23.6

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and N-(2-Methoxyethyl)ethylamine were converted to yield 4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl-ethyl-(2-methoxy-ethyl)-amine, MS: 403 (MH⁺, 1Cl).

23.7

According to the method in example 23, Methanesulfonic acid 4-[3-(4-chloro-phenyl)-benzo[d]isothiazol-6-yl]-butyl ester and Trimethylethylamine were converted to yield 6-(4-Azetidin-1-yl-butyl)-3-(4-chloro-phenyl)-benzo[d]isothiazole, MS: 357 (MH⁺, 1Cl).

Example 24

24.1

Method A: 75 mg (0.2 mmol) 2-((3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl)-ethyl-amino)-ethanol in 2 ml MeOH with 0.012 ml (0.2 mmol) AcOH were hydrogenated in the presence of 7 mg Pd/C 10% (16h), filtration over decalite and evaporation yielded 58 mg (66%) Z-2-((3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-allyl)-ethyl-amino)-ethanol acetate as light brown oil, MS: 373 (MH⁺, 1Cl).

24.2

Method B: 80 mg (0.25 mmol) {3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-dimethyl-amine in 2 ml EtOH were hydrogenated in the presence of 20 mg PtO₂. H₂O, purification by flash silica gel column (CH₂Cl₂/ MeOH 9:1) yielded 43 mg (53%) {3-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-propyl}-dimethyl-amine as a colorless oil, MS: 331 (MH⁺, 1Cl).

25 Example 25

25.1

10.09 g (70.5 mmol) 3-mercaptoanisole were added to a freshly prepared solution of 4.6 g (70.5 mmol) KOH in 75 ml ethanol and 30 ml water. Over a period of 30 min, 20.0 g (70.5 mmol) 2,4-dibromoacetophenone in 150 EtOAc were added at 0°C and the solution was stirred at RT over night. The suspension was concentrated and dissolved in EtOAc and water. The inorganic phase was extracted with EtOAc, the combined organic phases were washed with water and brine and dried over Na₂SO₄. Evaporation yielded 25.1g (quant) 1-(4-Bromo-phenyl)-2-(3-methoxy-phenylsulfanyl)-ethanone as white crystalline, MS: 336(M, 1Br). (In analogy to Jones, Charles D.; Jevnikar, Mary G.; Pike, Andrew J.; Peters,

- Mary K.; Black, Larry J.; Thompson, Allen R.; Falcone, Julie F.; Clemens, James A. Antiestrogens. 2. Structure-activity studies in a series of 3-aryl-2-arylbenzo[b]thiophene derivatives leading to [6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thien-3-yl]-[4-[2-(1-piperidinyl)ethoxy]phenyl] methanone hydrochloride (LY 156758), a remarkably effective
5 estrogen antagonist with only minimal intrinsic estrogenicity. J. Med. Chem. (1984), 27(8), 1057-66.)

25.2

- 60 g polyphosphoric acid were heated to 80°C. 11.85 g (35.14 mmol) 1-(4-Bromo-phenyl)-2-(3-methoxy-phenylsulfanyl)-ethanone were added in small portions, keeping the
10 temperature below 100°C. The reaction mixture was stirred at 85°C for 2h, cooled to 70°C and added to ice water. The aqueous phase was extracted with EtOAc three times, the organic phase was washed with brine and dried over Na₂SO₄. The solid was crysallized from EtOAc:MeOH to give 6.3 g (56%) 3-(4-Bromo-phenyl)-6-methoxy-benzo[b]thiophene as light yellow solid. MS: 318 (M, 1Br).

15 25.3

- 7.4 g (23.17 mmol) 3-(4-Bromo-phenyl)-6-methoxy-benzo[b]thiophene in 47 ml acetic acid were treated with 27 ml (62% aq) HBr at 125°C for 4h. The solution was concentrated in vacuo and the residue was dissolved in NaHCO₃ and EtOAc. Prior to extraction with EtOAc the pH was adjusted with 2M NaOH (pH 8). The combined organic phases were
20 washed with brine and dried with Na₂SO₄. Trituration with hexane gave 6.96 g (98%) 3-(4-Bromo-phenyl)-benzo[b]thiophen-6-ol as light green solid, MS: 304 (M, 1Br).

Example 26

26.1

- 14.27 ml (115.0 mmol) 3-mercaptoanisole were added to a solution of 7.3 g (115.0 mmol)
25 KOH in 120 ml ethanol and 50 ml water. 25.0 g (115.2 mmol) 4-fluorophenacyl bromide in 120 ml EtOAc were added slowly at 0°C and the solution was stirred at RT over night. The suspension was concentrated and dissolved in EtOAc and water. The inorganic phase was extracted with EtOAc, the combined organic phases were washed with water and brine and dried over Na₂SO₄. Evaporation yielded 33.2 g crude 1-(4-Fluoro-phenyl)-2-(3-methoxy-phenylsulfanyl)-ethanone as light yellow liquid.
30

26.2

- The crude product of experiment 26.1 was dissolved in toluene and evaporated three times prior to the reaction. To the resulting solid 305 ml (1.2 mol) BF₃·Et₂O were added slowly and the solution was stirred at RT over night. This solution was added slowly to NaOH,
35 and the pH was adjusted to pH 7. The solution was extracted with CH₂Cl₂, the organic

phase was washed with brine and dried over Na_2SO_4 , yielding 29.96 g crude 3-(4-Fluoro-phenyl)-6-methoxy-benzo[b]thiophene.

26.3

29.96 g 3-(4-Fluoro-phenyl)-6-methoxy-benzo[b]thiophene in 230 ml acetic acid were
5 treated with 130 ml (62% aq) HBr at 125°C for 4h. The solution was added slowly to a cooled solution of 210 g NaOH in 600 ml water, the pH was adjusted to 8-9, and the inorganic phase was extracted with EtOAc. The combined organic phases were washed with brine and dried with Na_2SO_4 . Column chromatography on silica gel with hexane: EtOAc 4:1 yielded 20.0 g (73%) 3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-ol as light
10 brown solid, MS: 244(M).

26.4

In analogy to examples 26.1 – 26.3, 3-mercaptoanisole and 2,4'-dibromopropiophenone were converted to yield 3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophen-6-ol as colorless amorphous solid, MS: 318 (M, 1Br).

15 26.5

In analogy to examples 26.1 – 26.3, 3-Mercaptoanisole and 2-Bromo-1-(4-trifluoromethyl-phenyl)-propan-1-one were converted to yield 2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-ol as colorless amorphous solid, MS: 308 (M).

Example 27

20 27.1

2.0 g (6.55 mmol) 3-(4-Bromo-phenyl)-benzo[b]thiophen-6-ol in 25 ml acetone were treated with 5.89 g (42.6 mmol, 6.5 eq) K_2CO_3 and 2.0 ml (16.38 mmol, 2.5 eq) 1,4-dibromobutane. The reaction mixture was stirred at 45°C for 10h, filtered and evaporated. The excess of dibromide was removed in vacuo to yield 2.7 g crude 6-(4-Bromo-butoxy)-
25 3-(4-bromo-phenyl)-benzo[b]thiophene, MS: 439 (M).

27.2

In analogy to example 27.1, 3-(4-Bromo-phenyl)-benzo[b]thiophen-6-ol and 1,3-dibromopropane were converted to yield 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene, MS: 424 (M, 2Br).

30 27.3

In analogy to example 27.1, 3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-ol and 1,3-dibromopropane were converted to yield 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene, MS: 364 (M, 1Br).

27.4

In analogy to example 27.1, 3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-ol and 1,4-dibromobutane were converted to yield 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene, MS: 378 (M, 1Br).

5 27.5

In analogy to example 27.1, 3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophen-6-ol and 1,4-dibromobutane were converted to yield 6-(4-bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene as colorless oil, MS: 452 (M, 2Br).

27.6

- 10 In analogy to example 27.1, 2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-ol and 1,4-dibromobutane were converted to yield 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene as colorless oil, MS: 443 (MH⁺, 1Br).

Example 28

28.1

- 15 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and N,N-dimethylamine in ethanol were converted to yield [4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl]-dimethyl-amine as light yellow semisolid, MS: 404 (MH⁺, 1Br).

28.2

- 20 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and N-Allylmethylamine were converted to yield Allyl-[4-[3-(4-bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl]-methyl-amine as light yellow oil, MS: 430 (MH⁺, 1Br).

28.3

- 25 In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene and diethanolamine were converted to yield 2-[[3-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-propyl]-(2-hydroxy-ethyl)-amino]-ethanol as light yellow oil, MS: 450 (MH⁺, 1Br).

28.4

- 30 In analogy to example 3.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene and bis(2-methoxyethyl)amine were converted to yield {3-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-bis-(2-methoxy-ethyl)-amine as light brown oil, MS: 478 (MH⁺, 1Br).

Example 29

1 eq of the corresponding bromide was treated with 3 eq of the corresponding amine in DMA (4-10 ml/mmol bromide) at RT until no starting material could be detected with TLC. The solution was concentrated and the residue was redissolved in CH₂Cl₂/ 5% aqueous NaHCO₃. The phases were separated, and the inorganic phase was extracted with CH₂Cl₂, the combined organic phases were washed with brine, and were dried over Na₂SO₄. The crude material was purified by flash chromatography. To yield the corresponding hydrochlorides, the free amines were treated with HCl in MeOH.

29.1

- 10 According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and 2-(methylamino)ethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amino)-ethanol-hydrochloride as light brown oil, MS: 434 (MH⁺, 1Br).

29.2

- 15 According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and N-(2-methoxyethyl)methylamine were converted to yield 4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl)-(2-methoxy-ethyl)-methylamine-hydrochloride as light brown semisolid, MS: 448 (MH⁺, 1Br).

29.3

- 20 According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and morpholine were converted to yield 4-[4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl]-morpholine-hydrochloride as light brown semisolid, MS: 446 (MH⁺, 1Br).

29.4

- 25 According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and 2-(ethylamino)ethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-ethanol-hydrochloride as orange oil, MS: 448 (MH⁺, 1Br).

29.5

- 30 According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and pyrrolidine were converted to yield 1-{4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-pyrrolidine-hydrochloride as grey semisolid, MS: 430 (MH⁺, 1Br).

29.6

According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and azetidine were converted to yield 1-{4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-azetidine-hydrochloride as light yellow oil, MS: 416
5 (MH⁺, 1Br).

29.7

According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and methylisopropylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-isopropyl-methyl-amine-hydrochloride as
10 white foam, MS: 432 (MH⁺, 1Br).

29.8

According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and diethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine-hydrochloride as orange oil, MS: 432
15 (MH⁺, 1Br).

29.9

According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and diethanolamine were converted to yield 2-[[4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol as light
20 yellow oil, MS: 464 (MH⁺, 1Br).

29.10

According to the method in example 29, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene and bis(2-methoxyethyl)amine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-bis-(2-methoxy-ethyl)-amine as light
25 brown oil, MS: 492 (MH⁺, 1Br).

Example 30

30.1

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and N-allylmethylamine were converted to yield Allyl-{3-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-methyl-amine as colorless oil, MS: 356 (MH⁺).
30

30.2

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and dimethylamine were converted to yield {3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-dimethyl-amine as colorless oil, MS: 330 (MH⁺).

30.3

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and 2-(methylamino)ethanol were converted to yield 2-({3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-methyl-amino)-ethanol as colorless oil, MS: 360 (MH⁺).

30.4

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and N-(2-methoxyethyl)methylamine were converted to yield {3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-(2-methoxy-ethyl)-methyl-amine as colorless oil, MS: 374 (MH⁺).

30.5

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and 2-(ethylamino)ethanol were converted to yield 2-(Ethyl-{3-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-amino)-ethanol as colorless oil, MS: 374 (MH⁺).

15 30.6

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and N-(2-methoxyethyl)ethylamine were converted to yield Ethyl-{3-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-(2-methoxy-ethyl)-amine as colorless oil, MS: 388 (MH⁺).

20 30.7

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and diethanolamine were converted to yield 2-[{3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-(2-hydroxy-ethyl)-amino]-ethanol as colorless oil, MS: 390 (MH⁺).

25 30.8

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and bis(2-methoxyethyl)amine were converted to yield {3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-bis-(2-methoxy-ethyl)-amine as colorless oil, MS: 418 (MH⁺).

30 30.9

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and pyrrolidine were converted to yield 1-{3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-pyrrolidine as colorless oil, MS: 356 (MH⁺).

30.10

In analogy to example 3.1, 6-(3-Bromo-propoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and azetidine were converted to yield 1-{3-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-propyl}-azetidine as colorless oil, MS: 342 (MH⁺).

5 30.11

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and N-allylmethylamine were converted to yield Allyl-{4-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine as colorless oil, MS: 370 (MH⁺).

30.12

10 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and dimethylamine were converted to yield {4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine as white semisolid, MS: 344 (MH⁺).

30.13

15 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and 2-(methylamino)ethanol were converted to yield 2-({4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amino)-ethanol as colorless oil, MS: 374 (MH⁺).

30.14

20 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and N-(2-methoxyethyl)methylamine were converted to yield {4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine as colorless oil, MS: 388 (MH⁺).

30.15

25 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and 2-(ethylamino)ethanol were converted to yield 2-(Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol as colorless oil, MS: 388 (MH⁺).

30.16

30 In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and N-(2-methoxyethyl)ethylamine were converted to yield Ethyl-{4-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-amine as colorless oil, MS: 402 (MH⁺).

30.17

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and diethanolamine were converted to yield 2-[{4-[3-(4-Fluoro-phenyl)-

benzo[b]thiophen-6-yloxy]-butyl)-(2-hydroxy-ethyl)-amino]-ethanol as white semisolid, MS: 404 (MH^+).

30.18

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and bis(2-methoxyethyl)amine were converted to yield {4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-bis-(2-methoxy-ethyl)-amine as colorless oil, MS: 432 (MH^+).

30.19

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and pyrrolidine were converted to yield 1-{4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-pyrrolidine as white semisolid, MS: 370 (MH^+).

30.20

In analogy to example 3.1, 6-(4-Bromo-butoxy)-3-(4-fluoro-phenyl)-benzo[b]thiophene and azetidine were converted to yield 1-{4-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-azetidine as colorless oil, MS: 356 (MH^+).

Example 31

A solution of 0.25 mmol (1 equivalent) of the corresponding bromide in 0.7 ml dry DMA was treated with a solution of 0.5 mmol (2 equivalents) of the corresponding secondary amine in 0.15 ml dry DMA at room temperature. After 16 h, additional 2 equivalents of secondary amine were added to the solution. The reaction mixture was allowed to stand over night at room temperature, treated with 0.2 ml formic acid and purified by preparative HPLC [RP-18, acetonitrile (0.1 % $HCOOH$)/water (0.1 % $HCOOH$), 10 % to 95 % acetonitrile]. After evaporation, the product was obtained as a mixture of amino hydrobromide and formate.

25 31.1

According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and Allyl-methyl-amine were converted to yield Allyl-{4-[3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine, MS: 444 (MH^+ , 1Br).

30 31.2

According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and Dimethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine, MS: 418 (MH^+ , 1Br).

31.3

According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and Ethyl-(2-hydroxy-ethyl)-amine were converted to yield 2-
({4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-
5 ethanol, MS: 462 (MH⁺, 1Br).

31.4

According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and Diethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine, MS: 446 (MH⁺, 1Br).

10 31.5

According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and piperidine were converted to yield 1-{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-piperidine, MS: 458 (MH⁺, 1Br).

31.6

15 According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and Azetidine were converted to yield 1-{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-azetidine, MS: 430 (MH⁺, 1Br).

31.7

According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-
20 methyl-benzo[b]thiophene and 2-(hydroxy-ethyl)-amino-ethanol were converted to yield 2-[[4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol, MS: 478 (MH⁺, 1Br).

31.8

According to the method in example 31, 6-(4-Bromo-butoxy)-2-methyl-3-(4-
25 trifluoromethyl-phenyl)-benzo[b]thiophene and 4-hydroxy-piperidine were converted to yield 1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-piperidin-4-ol as yellowish oil, MS: 464 (MH⁺).

31.9

According to the method in example 31, 6-(4-Bromo-butoxy)-2-methyl-3-(4-
30 trifluoromethyl-phenyl)-benzo[b]thiophene and N-propyl amine were converted to yield {4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-propyl-amine as yellowish oil, MS: 422 (MH⁺).

31.10

According to the method in example 31, 6-(4-Bromo-butoxy)-2-methyl-3-(4-

trifluoromethyl-phenyl)-benzo[b]thiophene and ethanolamine were converted to yield 2-[4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butylamino]-ethanol as colorless amorphous solid, MS: 424 (MH⁺).

31.11

- 5 According to the method in example 31, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene and 2-ethylamino-ethanol were converted to yield 2-(Ethyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol as yellowish oil, MS: 452 (MH⁺).

31.12

- 10 According to the method in example 31, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene and N-allyl-methylamine were converted to yield Allyl-methyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-amine as yellowish oil, MS: 434 (MH⁺).

31.13

- 15 According to the method in example 31, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene and piperidine were converted to yield 1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-butyl}-piperidine as yellowish oil, MS: 448 (MH⁺).

31.14

- 20 According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and N-(2-methoxy-ethyl)-ethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine as yellowish oil, MS: 476 (MH⁺, 1Br).

31.15

- 25 According to the method in example 31, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene and dimethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine as yellowish oil, MS: 418 (MH⁺, 1Br).

Example 32

32.1

30 1.3 g (2.95 mmol) 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene in 140 ml acetic acid were warmed to 50°C and 1.36 g (8.9 mmol) sodium perborate were added in portions. The reaction mixture was stirred at that temperature for 18 h, concentrated and dissolved in water, and extracted with EtOAc and CH₂Cl₂. The combined organic

phases were washed with brine and dried over Na_2SO_4 . Column chromatography on silica gel with EtOAc: hexane 1:4 yielded 802 mg (67%) 6-(4-Bromo-butoxy)-3-(4-bromophenyl)-benzo[b]thiophene 1,1-dioxide as light yellow solid, MS: 472 (MH^+ , 1Br).

32.2

- 5 In analogy to example 32.1, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene was converted to yield 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide as light yellow solid, MS: 458 (MH^+ , 1Br).

32.3

- 10 In analogy to example 32.1, 6-(4-bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene was converted to yield 6-(4-bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide as colorless solid, MS: 484 (M, 2Br).

32.4

- 15 In analogy to example 32.1, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene was converted to yield 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene 1,1-dioxide as colorless solid, MS: 475 (MH^+ , 1Br).

Example 33

- 1 eq of the corresponding bromide was treated with 3 eq of the corresponding amine in DMA (4-10 ml/mmol bromide) at RT until no starting material could be detected with
20 TLC. The solution was concentrated and the residue was redissolved in CH_2Cl_2 / 5% aqueous NaHCO_3 . The phases were separated, and the inorganic phase was extracted with CH_2Cl_2 , the combined organic phases were washed with brine, and were dried over Na_2SO_4 . The crude material was purified by flash chromatography. To yield the corresponding hydrochlorides, the free amines were treated with HCl in MeOH.

25 33.1

According to the method in example 33, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide and N-Allylmethylamine were converted to yield Allyl-{3-[3-(4-bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-propyl}-methylamine as light yellow solid, MS: 448(MH^+ , 1Br).

30 33.2

According to the method in example 33, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide and Dimethylamine were converted to yield {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-propyl}-dimethylamine as off-white solid, MS: 422 (MH^+ , 1Br).

33.3

According to the method in example 33, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide and N-Methylaminoethanol were converted to yield 2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-propyl}-methyl-amino)-ethanol as off-white semisolid, MS: 452 (MH⁺, 1Br).

33.4

According to the method in example 33, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide and N-Ethylaminoethanol were converted to yield 2-({3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-propyl}-ethyl-amino)-ethanol as light yellow semisolid, MS: 466 (MH⁺, 1Br).

33.5

According to the method in example 33, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide and N-(2-Methoxyethyl)methylamine were converted to yield {3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-propyl}-(2-methoxy-ethyl)-methyl-amine as off-white solid, MS: 466 (MH⁺, 1Br).

33.6

According to the method in example 33, 3-(4-Bromo-phenyl)-6-(3-bromo-propoxy)-benzo[b]thiophene 1,1-dioxide and azetidine were converted to yield 1-{3-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-propyl}-azetidine as off-white solid, MS: 434 (MH⁺, 1Br).

33.7

According to the method in example 33, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene 1,1-dioxide and N-Allylmethylamine were converted to yield Allyl-[4-[3-(4-bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl]-methyl-amine as off-white semisolid, MS: 462 (MH⁺, 1Br).

33.8

According to the method in example 33, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene 1,1-dioxide and Dimethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-dimethyl-amine as light yellow semisolid, MS: 436 (MH⁺, 1Br).

33.9

According to the method in example 33, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene 1,1-dioxide and N-Methylaminoethanol were converted to yield 2-({4-

[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amino)-ethanol as off-white solid, MS: 466 (MH^+ , 1Br).

33.10

According to the method in example 33, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene 1,1-dioxide and N-Ethylaminoethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-ethanol as off-white solid, MS: 480 (MH^+ , 1Br).

33.11

According to the method in example 33, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene 1,1-dioxide and N-(2-Methoxyethyl)methylamine were converted to yield {4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-(2-methoxy-ethyl)-methyl-amine as light yellow semisolid, MS: 480 (MH^+ , 1Br).

33.12

According to the method in example 33, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-benzo[b]thiophene 1,1-dioxide and azetidine were converted to yield 1-{4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-azetidine as off-white solid, MS: 434 (MH^+ , 1Br).

Example 34

A solution of 0.25 mmol (1 equivalent) of the corresponding bromide in 0.7 ml dry DMA was treated with a solution of 0.5 mmol (2 equivalents) of the corresponding secondary amine in 0.15 ml dry DMA at room temperature. After 16 h, additional 2 equivalents of secondary amine were added to the solution. The reaction mixture was allowed to stand over night at room temperature, treated with 0.2 ml formic acid and purified by preparative HPLC [RP-18, acetonitrile (0.1 % HCOOH)/water (0.1 % HCOOH), 10 % to 95 % acetonitrile]. After evaporation, the product was obtained as a mixture of amino hydrobromide and formate.

34.1

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and N-Methylallylamine were converted to yield Allyl-{4-[3-(4-bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine, MS: 476 (MH^+ , 1Br).

34.2

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and Diethylamine were converted to yield {4-[3-

(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl)-diethyl-amine, MS: 478 (MH⁺, 1Br).

34.3

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and N-Methylpropylamine were converted to yield
5 {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl)-methyl-propyl-amine, MS: 478 (MH⁺, 1Br).

34.4

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and Diethanolamine were converted to yield 2-[[4-
10 {3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl)-(2-hydroxy-ethyl)-amino]-ethanol, MS: 510 (MH⁺, 1Br).

34.5

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and 2-Ethylaminoethanol were converted to yield
15 2-({4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-amino)-ethanol, MS: 494 (MH⁺, 1Br).

34.6

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and N-(2-Methoxyethyl)ethylamine were
20 converted to yield {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-ethyl-(2-methoxy-ethyl)-amine, MS: 508 (MH⁺, 1Br).

34.7

According to the method in example 34, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene 1,1-dioxide and Trimethyleneamine were converted to yield 1-
25 {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-azetidine, MS: 462 (MH⁺, 1Br).

34.8

According to the method in example 34, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene 1,1-dioxide and 2-ethylamino-ethanol were
30 converted to yield 2-(Ethyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol as colorless oil, MS: 484 (MH⁺).

34.9

According to the method in example 34, 6-(4-Bromo-butoxy)-2-methyl-3-(4-

trifluoromethyl-phenyl)-benzo[b]thiophene 1,1-dioxide and diethylamine were converted to yield Diethyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-amine as yellowish solid, MS: 468 (MH⁺).

34.10

- 5 According to the method in example 34, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene 1,1-dioxide and N-methyl propyl-amine were converted to yield Methyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-propyl-amine as yellowish oil, MS: 468 (MH⁺).

34.11

- 10 According to the method in example 34, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene 1,1-dioxide and N-(2-methoxy-ethyl) ethyl-amine were converted to yield Ethyl-(2-methoxy-ethyl)-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-amine as yellowish oil, MS: 498 (MH⁺).

15 34.12

According to the method in example 34, 6-(4-Bromo-butoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene 1,1-dioxide and diethanol-amine were converted to yield 2-((2-Hydroxy-ethyl)-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-amino)-ethanol as colorless oil, MS:

20 500 (MH⁺).

Example 35

35.1

- To 7.0 g (15.6 mmol) 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol in 20 ml DMF were added 1.9 g (28.0 mmol, 1.8 eq) imidazole,
25 followed by 3.3 g (21.8 mmol, 1.4 eq) tert-butyldimethylchlorosilane in 20 ml DMF at 0°C. The solution was stirred at 50°C for 6 h, and at RT for 48 h. Additional 235 mg (1.5 mmol, 0.1 eq) tert-butyldimethylchlorosilane were added and the solution was stirred at 50°C for 2h. The solution was poured on an aqueous solution of NaHCO₃. The inorganic layer was extracted with ether, and the combined organic phases were washed with brine and dried
30 over Na₂SO₄. Column chromatography on silica gel with CH₂Cl₂:MeOH 95:5 gave 6.5 g (74%) {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine as brown oil, MS: 563 (MH⁺, 1Br).

35.2

Under argon a flask was charged with 26.0 mg (0.03 mmol, 0.05 eq)

tris(Dibenzylideneacetone) dipalladium, 16.9 mg (0.06 mmol, 0.1 eq) 2(di-tertbutylphosphino)biphenyl and 92.8 mg (0.97 mmol, 1.7 eq) sodium tert-butyrate, evacuated and backfilled with argon. 320 mg (0.57 mmol) 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine in 3 ml toluene were added, followed by a solution of 0.14 ml (1.4 mmol, 2.5 eq) piperidine in 3 ml toluene. The solution was heated to 80°C for 2d. The mixture was diluted with EtOAc and a saturated solution of Na₂CO₃ was added. The inorganic layer was extracted with EtOAc. The combined organic phases were washed with brine and dried over Na₂SO₄. Column chromatography yielded 248 mg (77 %) [2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-{4-[3-(4-piperidin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine as yellow oil, MS: 568 (MH⁺).

35.3

In analogy to example 35.2, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and Morpholine were converted to yield [2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-{4-[3-(4-morpholin-4-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine as yellow oil, MS: 570 (MH⁺).

35.4

In analogy to example 35.2, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and N-Methylaniline were converted to yield {4-[6-(4-{[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-methyl-phenyl-amine as yellow oil, MS: 590 (MH⁺).

35.5

In analogy to example 35.2, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and piperazine were converted to yield [2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-{4-[3-(4-piperazin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine as orange oil, MS: 569 (MH⁺).

35.6

In analogy to example 35.2, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and 1-acetylpiperazine were converted to yield 1-(4-{4-[6-(4-{[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-piperazin-1-yl)-ethanone as orange oil, MS: 611 (MH⁺).

35.7

In analogy to example 35.2, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-

[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and benzylamine were converted to yield Benzyl-{4-[6-(4-{[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-amine as orange oil, MS: 590 (MH⁺).

35.8

- 5 220 mg (0.39 mmol) [2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-{4-[3-(4-piperidin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine in 5 ml THF were treated with 0.6 ml 1M TBAF in THF for 1h at RT. The solution was concentrated and the crude product subjected to column chromatography on silica gel with CH₂Cl₂:MeOH 9:1 to give 146 mg (83%) 2-(Ethyl-{4-[3-(4-piperidin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol as yellow oil, MS: 454 (MH⁺).

35.9

- In analogy to example 35.8, [2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-{4-[3-(4-morpholin-4-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine were converted to yield 2-(Ethyl-{4-[3-(4-morpholin-4-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol as yellow oil, MS: 456 (MH⁺).

35.10

- In analogy to example 35.8, {4-[6-(4-{[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-methyl-phenyl-amine were converted to yield 2-[Ethyl-{4-[3-[4-(methyl-phenyl-amino)-phenyl]-benzo[d]isothiazol-6-yloxy]-butyl}-amino]-ethanol as yellow oil, MS: 476 (MH⁺).

35.11

- In analogy to example 35.8, [2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-{4-[3-(4-piperazin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine were converted to yield 2-(Ethyl-{4-[3-(4-piperazin-1-yl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol as colorless oil, MS: 455 (MH⁺).

35.12

- In analogy to example 35.8, 1-(4-{4-[6-(4-{[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-piperazin-1-yl)-ethanone were converted to yield 1-{4-[4-(6-{4-[Ethyl-(2-hydroxy-ethyl)-amino]-butoxy}-benzo[d]isothiazol-3-yl)-phenyl]-piperazin-1-yl}-ethanone as colorless oil, MS: 497 (MH⁺).

35.13

- In analogy to example 35.8, Benzyl-{4-[6-(4-{[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-amine were converted to yield 2-

({4-[3-(4-Benzylamino-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol as yellow oil, MS: 476 (MH⁺).

35.14

Under argon a flask was charged with 20.3 mg (0.02 mmol, 0.05 eq)
5 tris(dibenzylideneacetone) dipalladium, 13.2 mg (0.04 mmol, 0.1 eq) 2-(di-
tertbutylphosphino)biphenyl and 38.7 mg (0.89 mmol, 2 eq) sodium hydride (55% in
mineral oil), evacuated and backfilled with argon. 250 mg (0.44 mmol) 4-[3-(4-Bromo-
phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-
ethyl-amine in 4 ml toluene were added, followed by a solution of 0.031 ml (0.75 mmol,
10 1.7 eq) methanol in 2 ml toluene. The solution was heated to 80°C for 2d. The mixture was
diluted with EtOAc and a saturated solution of Na₂CO₃. The inorganic layer was extracted
with EtOAc. The combined organic phases were washed with brine and dried over Na₂SO₄.
Column chromatography on silica gel with CH₂Cl₂:MeOH 95:5 yielded 174 mg light
yellow oil which was dissolved in 5 ml THF and was treated with with 0.53 ml 1M Bu₄NF
15 in THF at RT for 1h. The solution was concentrated and the residue purified by column
chromatography to yield 2-[Ethyl-[4-(3-phenyl-benzo[d]isothiazol-6-yloxy)-butyl]-
amino]-ethanol as light yellow oil, MS: 371 (MH⁺).

35.15

Under argon a flask was charged with 10.0 mg (0.01 mmol, 0.03 eq)
20 tris(dibenzylideneacetone) dipalladium, 10 mg (0.02 mmol, 0.05 eq) R(+) 2,2'-
Bis(diphenylphosphino)-1,1'-Binaphthyl and 47.7 mg (0.5 mmol, 1.4 eq) sodium tert-
butylate, evacuated and backfilled with argon. 200 mg (0.35 mmol) 4-[3-(4-Bromo-
phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-
ethyl-amine in 4 ml toluene were added, followed by a solution of 0.071 ml (0.4mmol, 1.2
25 eq) benzophenone imine in 2 ml toluene. The solution was heated to 80°C over night. The
mixture was diluted with EtOAc and water. The inorganic layer was extracted with EtOAc.
The combined organic phases were washed with brine and dried over Na₂SO₄.

The residue was dissolved in 2 ml MeOH, 10 mg 10% Pd/C and 336 mg (5.3 mmol)
ammonium formate were added and the reaction mixture was stirred at 60°C for 3h.
30 Additional ammonium formate was added in portions until no starting material could be
detected. The solution was filtered over decalite and evaporated.

The crude product was dissolved in 1.5 ml THF and was treated with 0.24 ml 1M Bu₄NF in
THF at RT for 1h. The solution was concentrated and the residue purified by ion exchange
chromatography to yield 40 mg (29%, 3 steps) 2-({4-[3-(4-Amino-phenyl)-
35 benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol as orange oil, MS: 386(MH⁺).

Examples 36

36.1

- A flask was charged with 31.1 mg (0.04 mmol, 0.05 eq) bis(triphenylphosphine) palladium(II) chloride, evacuated and backfilled with argon. 500 mg (0.89 mmol) 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and 7.0 mg (0.03 mmol, 0.03 eq) triphenylphosphine in 5 ml piperidine were added, and the solution stirred for 20 min. 1.7 mg (0.01 mmol, 0.01 eq) CuI were added, followed by 113 μ l (1.06 mmol, 1.2 eq) 1-Dimethylamino-2-propyne. The solution was stirred at 50°C over night and at 70°C for 1h. The mixture was poured on water and EtOAc, the layers were separated and the inorganic layer was extracted with EtOAc. The combined organic phases were washed with brine and dried over Na₂SO₄. Column chromatography on silica gel with CH₂Cl₂:MeOH 95:5 yielded 350 mg (70%) (3-{4-[6-(4-{[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-prop-2-ynyl)-dimethyl-amine as brown oil.
- 350 mg (0.62 mmol) (3-{4-[6-(4-{[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amino}-butoxy)-benzo[d]isothiazol-3-yl]-phenyl}-prop-2-ynyl)-dimethyl-amine in 5 ml THF were treated with 0.92 ml 1M Bu₄NF in THF at RT for 1h. The solution was concentrated and the residue was purified by column chromatography on silica gel to give 185 mg (66%) 2-[(4-{3-[4-(3-Dimethylamino-prop-1-ynyl)-phenyl]-benzo[d]isothiazol-6-yloxy}-butyl)-ethyl-amino]-ethanol as light brown oil, MS: 452 (MH⁺).

36.2

- In analogy to example 36.1, {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and 2-Propyn-1-ol were converted to yield 3-[4-(6-{4-[Ethyl-(2-hydroxy-ethyl)-amino]-butoxy}-benzo[d]isothiazol-3-yl)-phenyl]-prop-2-yn-1-ol as yellow oil, MS: 425 (MH⁺).

36.3

- In analogy to example 36.1, {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-ethyl-amine and N-Methylpropargylamine were converted to yield 2-[Ethyl-(4-{3-[4-(3-methylamino-prop-1-ynyl)-phenyl]-benzo[d]isothiazol-6-yloxy}-butyl)-amino]-ethanol as brown oil, MS: 438 (MH⁺).

Example 37

37.1

- To 14.3 ml (146.3 mmol) 2-Ethylaminoethanol in 60 ml THF were added 362 mg (3.7 mmol) CuCl, followed by 7.5 g (73.1 mmol) 3-Chloro-3-methyl-1-butyne. The solution

was stirred at RT for 2h, the solvent was removed and the residue dissolved in ether/2M HCl. The phases were separated and after addition of Na_2CO_3 to the aqueous phase (pH9), it was extracted with ether. The ether phase was dried over Na_2SO_4 and concentrated. Distillation by Kugelrohr yielded 2-[(1,1-Dimethyl-prop-2-ynyl)-ethyl-amino]-ethanol as
5 colorless oil, MS: 155 (M).

37.2

In analogy to example 37.1, 3-Chloro-3-methyl-1-butyne and Allylmethylamine were converted to yield Allyl-(1,1-dimethyl-prop-2-ynyl)-methyl-amine as white solid, MS: 137(M).

10 37.3

In analogy to example 37.1, 3-Chloro-3-methyl-1-butyne and N-(2-Methoxyethyl)methylamine were converted to yield (1,1-Dimethyl-prop-2-ynyl)-(2-methoxy-ethyl)-methyl-amine as colorless oil, MS: 155 (M).

37.4

15 In analogy to example 37.1, 3-Chloro-3-methyl-1-butyne and N-(2-Methoxyethyl)ethylamine were converted to yield (1,1-Dimethyl-prop-2-ynyl)-ethyl-(2-methoxy-ethyl)-amine as colorless oil, MS: 170 (MH^+).

37.5

According to Ishihara, Kazuaki; Kubota, Manabu; Kurihara, Hideki; Yamamoto, Hisashi.
20 Scandium Trifluoromethanesulfonate as an Extremely Active Lewis Acid Catalyst in Acylation of Alcohols with Acid Anhydrides and Mixed Anhydrides. J. Org. Chem. (1996), 61(14), 4560-4567:

To 3.1 g (25 mmol) 1-ethynyl-1-cyclohexanol in 25 ml acetonitrile were added 7.1 ml (75 mmol, 3 eq) acetic anhydride at -20°C . 12.3 mg (0.03 mmol) scandiumtriflate in 300 μl
25 THF were added and the solution was stirred for 2h. Additional 12.3 mg (0.03 mmol) scandiumtriflate in 300 μl THF were added and stirring was continued for 30 min. A saturated aqueous solution of NaHCO_3 and ether were added. The layers were separated and the inorganic one was extracted with ether. The combined organic phases were washed with brine and dried over Na_2SO_4 . Evaporation yielded 4.6g crude 1-ethynylcyclohexyl
30 acetate as colorless oil, MS: 166 (M).

37.6

Analogously to Imada, Yasushi; Yuasa, Mari; Nakamura, Ishin; Murahashi, Shun-Ichi. Copper(I)-Catalyzed Amination of Propargyl Esters. Selective Synthesis of

Propargylamines, 1-Alken-3-ylamines, and (Z)-Allylamines. J. Org. Chem. (1994), 59(9), 2282-4.

To 1.17 ml (12 mmo) ethylaminoethanol in 8 ml THF were added 29.8 mg (0.3 mmol) CuCl, followed by 1g (6.02 mmol) 1-ethynylcyclohexyl acetate. The solution was stirred at
5 RT for 3h, the solvent was removed and the residue dissolved in ether/2M HCl. The phases were separated, and after addition of NaOH (pH 11) to the aqueous phase, it was extracted with ether. The ether phase was dried over Na₂SO₄ and concentrated. Distillation by Kugelrohr yielded 2-[Ethyl-(1-ethynyl-cyclohexyl)-amino]-ethanol as colorless oil, MS: 196 (MH⁺).

10 37.7

In analogy to example 37.5 and 37.6, 3-Butyn-2-ol and 2-Ethylaminoethanol were converted to yield 2-[Ethyl-(1-methyl-prop-2-ynyl)-amino]-ethanol as colorless oil, MS: 141 (M).

37.8

15 In analogy to example 37.5 and 37.6, 3-Butyn-2-ol and N-(Methoxyethyl)methylamine were converted to yield (2-Methoxy-ethyl)-methyl-(1-methyl-prop-2-ynyl)-amine as colorless oil, MS: 141 (M).

37.9

In analogy to example 37.5 and 37.6, 3-Butyn-2-ol and N-(Methoxyethyl)ethylamine were
20 converted to yield Ethyl-(2-methoxy-ethyl)-(1-methyl-prop-2-ynyl)-amine as colorless oil, MS: 155 (M).

37.10

In analogy to example 37.5 and 37.6, 1-Ethynylcyclopentanol and N-Methyl-allylamine were converted to yield Allyl-(1-ethynyl-cyclopentyl)-methyl-amine as colorless oil, MS:
25 163 (M).

Example 38

38.1

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and Allyl-(1,1-dimethyl-prop-2-ynyl)-methyl-amine were
30 converted to yield Allyl-{1,1-dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-methyl-amine as brown oil, MS: 415 (MH⁺).

38.2

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-

benzo[d]isothiazol-6-yl ester and 2-[(1,1-Dimethyl-prop-2-ynyl)-ethyl-amino]-ethanol were converted to yield 2-((1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl)-ethyl-amino)-ethanol as off-white foam, MS: 433 (MH⁺).

5 38.3

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and (1,1-Dimethyl-prop-2-ynyl)-(2-methoxy-ethyl)-methyl-amine were converted to yield {1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-(2-methoxy-ethyl)-methyl-amine as brown oil,
10 MS: 433 (MH⁺).

38.4

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and (1,1-Dimethyl-prop-2-ynyl)-ethyl-(2-methoxy-ethyl)-amine were converted to yield {1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-ethyl-(2-methoxy-ethyl)-amine as brown oil, MS:
15 447 (MH⁺).

38.5

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and 2-[Ethyl-(1-ethynyl-cyclohexyl)-amino]-ethanol were
20 converted to yield 2-(Ethyl-{1-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ylethynyl]-cyclohexyl}-amino)-ethanol as brown oil, MS: 473.3 (MH⁺).

38.6

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and 2-[Ethyl-(1-methyl-prop-2-ynyl)-amino]-ethanol were
25 converted to yield 2-(Ethyl-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol as brown oil, MS: 419 (MH⁺).

38.7

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and (2-Methoxy-ethyl)-methyl-(1-methyl-prop-2-ynyl)-
30 amine were converted to yield (2-Methoxy-ethyl)-methyl-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine as brown oil, MS: 419 (MH⁺).

38.8

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-

benzo[d]isothiazol-6-yl ester and Ethyl-(2-methoxy-ethyl)-(1-methyl-prop-2-ynyl)-amine were converted to yield Ethyl-(2-methoxy-ethyl)-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amine as brown oil, MS: 433 (MH⁺).

38.9

- 5 In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl ester and Allyl-(1-ethynyl-cyclopentyl)-methyl-amine were converted to yield Allyl-methyl-{1-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-ylethynyl]-cyclopentyl}-amine as light yellow oil, MS: 441 (MH⁺).

Example 39

10 39.1

To 540 mg (11.37 mmol) 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol in 6 ml CH₂Cl₂ were added 0.22 ml (1.7 mmol) DAST at -78°C. After stirring at that temperature for 3.5 h, additional 0.2 ml (1.6 mmol) DAST were added and the mixture was slowly warmed to RT over night. The solution was added to a
15 cooled aqueous solution of Na₂CO₃ and extracted with EtOAc. The organic phase was washed with brine and dried over Na₂SO₄. Column chromatography CH₂Cl₂:MeOH 95:5 yielded 79.3 mg (16%) Ethyl-(2-fluoro-ethyl)-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine as brown oil, MS: 441 (MH⁺).

39.2

- 20 In analogy to example 39.1, 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-amino)-ethanol was converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-fluoro-ethyl)-amine as light yellow oil, MS: 451 (MH⁺, 1Br).

Example 40:

25 40.1

- 200 mg (0.54 mmol) 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-2-methyl-but-3-yn-2-ol in 2 ml THF were treated with 36 µl (5.6 mmol) 1.6 M N-Butyllithium in hexane at -78°C for 30 min. 112 µl (0.81 mmol, 1.5 eq) Triethylamine in 0.5 ml THF and 117 µl (0.8 mmol, 1.5 eq) Diethylchlorophosphate in 0.5 ml THF were added and the solution was
30 stirred at -78°C over night and at 0°C for 4h. Water was added, and the inorganic phase was extracted with EtOAc, the combined organic phases were washed with brine and dried over Na₂SO₄. Column chromatography on silica gel with hexane:EtOAc 3:1 gave 90 mg (33%) Phosphoric acid 3-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-1,1-dimethyl-prop-2-ynyl ester diethyl ester as yellow semisolid, MS: 508 (MH⁺, 1Br).

40.2

To 45 mg (0.09 mmol) Phosphoric acid 3-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-1,1-dimethyl-prop-2-ynyl ester diethyl ester and 16 μ l (0.18 mmol) 2-Ethylaminoethanol in 0.5 ml THF 10.2 mg (0.01 mmol) tetrakis(triphenylphosphine)palladium was added and
5 the suspension stirred at 50°C over night. The mixture was added to a saturated aqueous solution of NaHCO₃ and EtOAc, the phases were separated and the inorganic one was extracted with EtOAc. The combined organic phases were washed with brine and dried over Na₂SO₄. Column chromatography on silica gel with CH₂Cl₂:MeOH 95:5 gave 2-((3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-1,1-dimethyl-prop-2-ynyl)-ethyl-amino)-
10 ethanol as yellow semisolid, MS: 443 (MH⁺, 1Br).

Example 41:

41.1

Under argon a flask was charged with 1.3 g (2.9 mmol) 2-((4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl)-ethyl-amino)-ethanol in 10 ml piperidine and 170 mg
15 (0.14 mmol, 0.05 eq) Pd(PPh₃)₄ and 28 mg (0.14 mmol, 0.05 eq) CuI. After stirring for 10 min, 1 ml (4.44 mmol, 1.5 eq) (triisopropylsilyl)acetylene was added over a period of 1h to this solution. The solution was stirred at 80°C for 1.5h. The solution was poured on ice water, acidified and the inorganic layer was extracted with EtOAc. The combined organic phases were washed with brine and dried over Na₂SO₄.

20 The crude product was dissolved in 20 ml THF and treated with 3.75 ml 1M Bu₄NF in THF at RT for 1h. The solution was diluted with EtOAc and a solution of Na₂CO₃. The inorganic layer was extracted with EtOAc, the combined organic phases were washed with brine and dried over Na₂SO₄. Column chromatography on silica with CH₂Cl₂:MeOH 9:1 yielded 160 mg (19%) 2-(Ethyl-{4-[3-(4-ethynyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol as brown oil, MS: 395 (MH⁺).
25

41.2

150 mg (0.4 mmol) 2-(Ethyl-{4-[3-(4-ethynyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol in 6 ml EtOAc were hydrogenated in the presence of 50 mg 10% Pd/C, filtration over decalite, followed by purification by preparative HPLC yielded 2-
30 (Ethyl-{4-[3-(4-ethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-ethanol as light yellow oil, MS: 399 (MH⁺).

Example 42

42.1

To a suspension of 0.74 g (16.9 mmol) NaH (55% in mineral oil) in 40 ml THF was added

a solution of 4 g (13.1 mmol) 3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-ol in 20 ml THF at RT. The solution was stirred at RT for 1h, 2.1 ml (14.3 mmol) tbutyl bromoacetate was added. The solution was stirred at RT over night, 1M KHSO₄ was added carefully and the layers separated. The inorganic one was extracted with EtOAc, the combined organic
5 phases were washed with water and brine and dried over Na₂SO₄. Column chromatography on silica gel with EtOAc: hexane 1:4 yielded 4.95 g (90%) [3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-acetic acid tert-butyl ester as colorless oil, MS: 420 (MH⁺, 1Br).

42.2

10 3.89 g (9.3 mmol) [3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-acetic acid tert-butyl ester in 35 ml CH₂Cl₂ were added 17 ml TFA at 0°C. The solution was stirred at RT for 2h, concentrated and titrated with ether to yield 3 g (89%) [3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-acetic acid as white solid, MS: 362(M-H, 1Br).

42.3

15 To 400 mg (1.1 mmol) trans-5-[4-(tert-Butoxycarbonyl-methyl-amino)-cyclohexyl]-pentanoic acid in 10 ml CH₂Cl₂ were added 0.82 ml (1.6 mmol, 1.5 eq) 1M dimethylamine in THF and 0.18 ml (1.6 mmol, 1.5 eq) NMM. The solution was cooled to 0°C and 274 mg (1.43 mmol, 1.3 eq) EDCI and 30 mg (0.2 mmol) HOBT were added. The mixture was stirred at RT over night, partitioned between CH₂Cl₂ and a saturated aqueous solution of
20 NaHCO₃. The organic phase was washed with KHSO₄ and brine, dried over Na₂SO₄ and evaporated. Column chromatography with EtOAc/hexane 2:1 gave 370 mg (86%) 2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-N,N-dimethyl-acetamide as white solid, mp 168°C, MS: 391 (MH⁺, 1Br).

42.4

25 Analogously to example 42.3, from [3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-acetic acid and N,O-Dimethylhydroxylamine hydrochloride was prepared 2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-N-methoxy-N-methyl-acetamide as white solid, mp 159°C, MS: 407 (MH⁺, 1Br).

42.5

30 To a solution of 160 mg (0.41 mmol) 2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-N,N-dimethyl-acetamide in 4 ml THF was added 95 mg (4.1 mmol, 1 eq) zirconium(IV) chloride at -10°C and the reaction mixture was stirred for additional 30min. 0.82 ml 3M (24.5 mmol, 6 eq) methylmagnesium bromide in THF were added at -10°C and the mixture was warmed to RT. After stirring for 1h, the mixture was added to a
35 mixture of 30% NaOH and CH₂Cl₂. The inorganic phase was extracted with CH₂Cl₂ and the combined organic phases were washed with brine and dried over Na₂SO₄ and

evaporated. The residue was purified by column chromatography on silica gel with CH₂Cl₂ to CH₂Cl₂:MeOH 9:1 to yield 63 mg (38%) {2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-1,1-dimethyl-ethyl}-dimethyl-amine as colorless oil, MS: 405 (MH⁺, 1Br).

42.6

- 5 To 208 mg (0.51 mmol) 2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-N-methoxy-N-methyl-acetamide in 8 ml THF were added 0.51 ml 3M (1.53 mmol, 3 eq) methylmagnesiumbromide in THF at -75°C. The solution was slowly warmed to RT over night, a saturated aqueous solution of NH₄Cl was added and the mixture stirred for 30 min. The phases were separated and the inorganic phase was extracted with EtOAc. The
10 combined organic phases were dried over Na₂SO₄ and evaporated. Column chromatography on silica gel with EtOAc:hexane 1:4 yielded 109 mg (53%) 1-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propan-2-one as white solid, MS: 361 (M, 1Br).

42.7

- 15 74 mg (0.20 mmol) 1-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propan-2-one were treated with 133 µM 2M(0.17 mmol) dimethylamine in THF and 60 µM (0.20 mmol) tetraisopropyl orthotitanate. The solution was stirred at RT for 3.5 h. The mixture was diluted with 2 ml ethanol and 15 mg (0.2 mmol) NaCNBH₃ were added, and stirring was continued over night. NaHCO₃ and EtOAc were added, the inorganic phase extracted with
20 EtOAc. The organic phases were washed with water and brine, dried over Na₂SO₄ and evaporated. Column chromatography with CH₂Cl₂:MeOH 9:1 gave {2-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-1-methyl-ethyl}-dimethyl-amine as colorless oil, MS: 391 (MH⁺, 1Br).

Example 43

25 43.1

In analogy to example 12.1, 2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-ol (example 26.5) was treated with trifluoromethane sulfonic anhydride in pyridine to yield the Trifluoro-methanesulfonic acid 2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl ester as light brown oil, MS: 440 (M⁺).

30 43.2

In analogy to example 12.1, 3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-ol and trifluoromethane sulfonic anhydride were converted to yield Trifluoro-methanesulfonic acid 3-(4-fluoro-phenyl)-benzo[b]thiophene-6-yl ester as brown oil, MS: 376 (M).

Example 44**44.1**

In analogy to example 13.1, the Trifluoro-methanesulfonic acid 2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl ester was reacted with 2-propyn-1-ol to
5 yield the 3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-yn-1-ol as yellow oil, MS: 346 (M^+).

44.2

In analogy to example 13.1, the Trifluoro-methanesulfonic acid 2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl ester was reacted with 3-butyne-1-ol to
10 yield the 4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-but-3-yn-1-ol as brown oil, MS: 360 (M^+).

44.3

In analogy to example 13.1, the Trifluoro-methanesulfonic acid 2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl ester was reacted with 4-pentyne-1-ol to
15 yield the 5-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol as brown oil, MS: 374 (M^+).

44.4

In analogy to example 14.1, Trifluoro-methanesulfonic acid 3-(4-fluoro-phenyl)-benzo[b]thiophene-6-yl ester and 4-pentyne-1-ol were converted to yield 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol as light green solid, MS: 310 (M).

Example 45**45.1**

In analogy to example 20.1, the 3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-yn-1-ol was hydrogenated to yield the 3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propan-1-ol as colorless oil, MS: 350 (M^+).

45.2

In analogy to example 20.1, the 4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-but-3-yn-1-ol was hydrogenated to yield the 4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butan-1-ol as yellowish oil, MS: 364 (M^+).

Example 46

46.1

In analogy to example 21.1, the 3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propan-1-ol was treated with methanesulfonic acid chloride to
5 yield the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester as yellow oil, MS: 428 (M^+).

46.2

In analogy to example 21.1, the 4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butan-1-ol was treated with methanesulfonic acid chloride to
10 yield the Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester as yellowish oil, MS: 442 (M^+).

46.3

In analogy to example 21.1, the 5-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol was treated with methanesulfonic acid chloride to
15 yield the Methanesulfonic acid 5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl ester as yellow oil, MS: 452 (M^+).

46.4

In analogy to example 21.1, the 3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-yn-1-ol was treated with methanesulfonic acid chloride to
20 yield a mixture of the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl ester and the 6-(3-Chloro-prop-1-ynyl)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene as yellow oil, used without further characterization.

Example 47

25 47.1

In analogy to example 32.1, the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester was converted to yield the Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-propyl ester as yellow oil, MS: 478 (MNH_4^+).

30 47.2

In analogy to example 32.1, the Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester was converted to yield the Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl ester as yellowish oil, MS: 474 (M^+).

47.3

In analogy to example 32.1, the Methanesulfonic acid 5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl ester was converted to yield the Methanesulfonic acid 5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-
5 benzo[b]thiophen-6-yl]-pent-4-ynyl ester as colorless foam, MS: 484 (M^+).

Example 48

In analogy to the method described in example 33, Methanesulfonic acid esters or Chlorides were treated with secondary or primary amines in N,N-dimethylacetamide between room temperature and 60°C to yield tertiary or secondary amine products as
 5 listed in the following table. The compounds were obtained as colorless, yellowish or brown oils.

Example	Compound	MS MH ⁺	Methanesulfonate /Chloride	Amine
48.1	Methyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-propyl-amine	402	mixture of the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl ester and the 6-(3-Chloro-prop-1-ynyl)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene	N-methyl-propylamine
48.2	2-(Ethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-amino)-ethanol	418	mixture of the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl ester and the 6-(3-Chloro-prop-1-ynyl)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene	2-ethylamino-ethanol
48.3	Diethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-amine	402	mixture of the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl ester and the 6-(3-Chloro-prop-1-ynyl)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene	diethyl-amine

48.4	Allyl-methyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl}-amine	400	mixture of the Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-prop-2-ynyl ester and the 6-(3-Chloro-prop-1-ynyl)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene	N-allyl-methyl-amine
48.5	Allyl-methyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amine	428	Methanesulfonic acid 5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	N-allyl-methyl-amine
48.6	Methyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-propyl-amine	430	Methanesulfonic acid 5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	N-methyl-propyl-amine
48.7	2-(Ethyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amino)-ethanol	446	Methanesulfonic acid 5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	2-ethylamino-ethanol
48.8	Diethyl-{5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amine	430	Methanesulfonic acid 5-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	diethyl-amine
48.9	1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-azetidine	390	Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester	azetidine

48.10	1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-piperidin-4-ol	434	Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester	4-hydroxy-piperidine
48.11	1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-piperidine	418	Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester	piperidine
48.12	1-{3-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-pyrrolidine	404	Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester	pyrrolidine
48.13	2-(Ethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-amino)-ethanol	422	Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester	2-ethylamino-ethanol
48.14	Allyl-methyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-amine	404	Methanesulfonic acid 3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl ester	N-allyl-methylamine
48.15	Allyl-methyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-amine	418	Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester	N-allyl-methylamine
48.16	2-(Ethyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-amino)-ethanol	436	Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester	2-ethylamino-ethanol
48.17	2-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butylamino}-ethanol	408	Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester	ethanolamine

48.18	1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-piperidine	432	Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester	piperidine
48.19	1-{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-piperidin-4-ol	448	Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester	4-hydroxy-piperidine
48.20	{4-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-propyl-amine	406	Methanesulfonic acid 4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl ester	propylamine
48.21	2-(Ethyl-{3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl}-amino)-ethanol	454	Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl ester	2-ethylamino-ethanol
48.22	2-{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propylamino}-ethanol	426	Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl ester	ethanolamine
48.23	{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl}-propyl-amine	424	Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl ester	N-propyl-amine
48.24	1-{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl}-piperidin-4-ol	466	Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl ester	4-hydroxy-piperidine

48.25	1-{3-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl}-piperidine	450	Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl ester	piperidine
48.26	Allyl-methyl-{3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl}-amine	436	Methanesulfonic acid 3-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-propyl ester	N-allyl-methylamine
48.27	2-(Ethyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl}-amino)-ethanol	468	Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl ester	2-ethylamino-ethanol
48.28	Allyl-methyl-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl}-amine	450	Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl ester	N-allyl-methylamine
48.29	1-{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl}-piperidine	464	Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl ester	piperidine
48.30	1-{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl}-piperidin-4-ol	480	Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-11 6-benzo[b]thiophen-6-yl]-butyl ester	4-hydroxy-piperidine

48.31	{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl}-propyl-amine	438	Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl ester	N-propyl-amine
48.32	2-{4-[2-Methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butylamino}-ethanol	440	Methanesulfonic acid 4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-butyl ester	ethanolamine
48.33	2-(Ethyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amino)-ethanol	478	Methanesulfonic acid 5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	2-ethylamino-ethanol
48.34	Methyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-propyl-amine	462	Methanesulfonic acid 5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	N-methyl-propylamine
48.35	Allyl-methyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amine	460	Methanesulfonic acid 5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	N-allyl-methylamine
48.36	Diethyl-{5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amine	462	Methanesulfonic acid 5-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yl]-pent-4-ynyl ester	diethyl-amine

Example 49

49.1

In analogy to example 19.1, 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol and 2-(Methylamino)ethanol were converted to yield 2-({5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-methyl-amino)-ethanol as light brown oil, MS: 368 (MH⁺).

49.2

In analogy to example 19.1, 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol and 2-(Ethylamino)ethanol were converted to yield 2-(Ethyl-{5-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-amino)-ethanol as light brown oil, MS: 382 (MH⁺).

49.3

RO0721678-000: In analogy to example 19.1, 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol and Dimethylamine were converted to yield {5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-dimethyl-amine as light brown semisolid, MS: 338 (MH⁺).

49.4

In analogy to example 19.1, 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol and N-Allylmethylamine were converted to yield Allyl-{5-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-methyl-amine as brown oil, MS: 364 (MH⁺).

49.5

In analogy to example 19.1, 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol and N-(Methoxyethyl)ethylamine were converted to yield Ethyl-{5-[3-(4-fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-(2-methoxy-ethyl)-amine as light brown oil, MS: 396 (MH⁺).

49.6

In analogy to example 19.1, 5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-yn-1-ol and N-(Methoxyethyl)methylamine CH₂Cl₂ were converted to yield {5-[3-(4-Fluoro-phenyl)-benzo[b]thiophen-6-yl]-pent-4-ynyl}-(2-methoxy-ethyl)-methyl-amine as brown oil, MS: 382 (MH⁺).

49.7

In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and N-Methylallylamine were converted to yield Allyl-{4-[3-(4-bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-methyl-amine as colorless oil, MS: 415 (MH⁺, 1Br).

49.8

In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and Dimethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-dimethyl-amine as brown oil, MS: 389 (MH⁺, 1Br).

5 49.9

In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and 2-(Methylamino)ethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-methyl-amino)-ethanol as brown oil, MS: 419 (MH⁺, 1Br).

49.10

10 In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and 2-(Ethylamino)ethanol were converted to yield 2-({4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-amino)-ethanol as yellow oil, MS: 433 (MH⁺, 1Br).

49.11

15 In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and N-(Methoxyethyl)ethylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-(2-methoxy-ethyl)-amine as yellow oil, MS: 447 (MH⁺, 1Br).

49.12

20 In analogy to example 19.1, 4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butan-1-ol and N-(Methoxyethyl)methylamine were converted to yield {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-(2-methoxy-ethyl)-methyl-amine as yellow oil, MS: 433 (MH⁺, 1Br).

Example 50

50.1

25 4.0 g (13.0 mmol) 2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-ol (example 26.5) dissolved in 30 ml N,N-dimethylformamide were treated with 2.69 g (19.5 mmol, 1.5 eq) K₂CO₃ and 3.06 ml (19.5 mmol, 1.5 eq) 2-(2-Bromo-ethoxy)-tetrahydro-pyran. The reaction mixture was stirred at 120°C for 3 hours, cooled to room temperature, poured into 100 ml of ice-water and extracted 3 times with 50 ml of ether. The combined
30 ether phases were washed with brine, dried over magnesium sulfate and evaporated under reduced pressure. The residue formed was filtered over silica gel with dichloromethane as the eluent giving 5.76 g crude 2-{2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethoxy}-tetrahydro-pyran. This crude product was
35 subsequently dissolved in 70 ml of 2N HCl in methanol and the reaction mixture stirred for 1 hour at room temperature. Subsequently, it was poured into a diluted sodium

hydroxide/ice solution and extracted 3 times with 150 ml of dichloromethane. The combined dichloromethane phases were washed with brine, dried over magnesium sulfate and evaporated under reduced pressure. The residue formed was chromatographed on silica gel with dichloromethane as the eluent giving 4.3 g (93%) 2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethanol as colorless solid, MS: 352 (M^+).

50.2

In analogy to example 21.1, 2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethanol and methanesulfonic acid chloride were converted to yield Methanesulfonic acid 2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl ester as colorless solid, MS: 430 (M^+).

50.3

0.300 g (0.697 mmol) Methanesulfonic acid 2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl ester dissolved in 2.0 ml of N,N-dimethylformamide were treated with 0.137 g (2.09 mmol) sodium azide and the reaction mixture heated to 80°C for 1 hour. It was then cooled to room temperature, poured into 30 ml of ice-water and extracted 3 times with 10 ml of ether. The combined ether phases were washed with brine, dried over magnesium sulfate and evaporated under reduced pressure giving 0.262 g (99.6%) 6-(2-Azido-ethoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene as colorless oil, MS: 377 (M^+).

50.4

0.230 g (0.610 mmol) 6-(2-Azido-ethoxy)-2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophene and 0.484 g (1.83 mmol) of triphenylphosphine were dissolved in 1.6 ml of tetrahydrofuran. After stirring of the reaction mixture for 15 minutes at room temperature, 0.1 ml of water were added and the reaction mixture heated to 60°C for 16 hours. It was then evaporated under reduced pressure, poured into 50 ml of water and extracted 3 times with 10 ml of dichloromethane. The combined dichloromethane phases were washed with brine, dried over magnesium sulfate and evaporated under reduced pressure. The residue formed was chromatographed on silica gel with a 9:1 v/v mixture of dichloromethane and methanol as the eluent giving 0.200 g (93.4%) 2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethylamine as colorless oil, MS: 352 (MH^+).

50.5

In analogy to the procedures described in examples 49.3 and 49.4, 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene (example 27.5) was reacted with sodium azide to yield 6-(4-Azido-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene,

which was further reduced with triphenylphosphine and water to yield 4-[3-(4-Bromophenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butylamine as colorless oil, MS: 390 (MH^+ , 1Br).

Example 51

5 51.1

In analogy to the method described in example 17.1, Methanesulfonic acid 2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl ester (example 50.2) was treated with methylamine in ethanol and N,N-dimethylacetamide to yield Methyl-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine as
10 colorless oil, MS: 366 (MH^+).

51.2

In analogy to the method described in example 17.1, 6-(4-Bromo-butoxy)-3-(4-bromophenyl)-2-methyl-benzo[b]thiophene (example 27.5) was treated with methylamine in ethanol and N,N-dimethylacetamide to yield {4-[3-(4-Bromo-phenyl)-2-methyl-
15 benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine as colorless oil, MS: 405 (MH^+).

Example 52

52.1

A solution of 40 mg (0.109 mmol) Methyl-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine (example 51.1), 0.0211 g (0.164 mmol) of 4-
20 chloro-2-methyl-pyrimidine [Ger. Offen. (1990), DE3905364 A1] and 0.038 ml (0.22 mmol) N-ethyl-diisopropylamine in 1 ml of N,N-dimethylformamide was stirred for 2 hours at 80°C. The reaction mixture was then cooled to room temperature, poured into 30 ml of ice-water and extracted 3 times with 10 ml of ether. The combined ether phases were washed with brine, dried over magnesium sulfate and evaporated under reduced pressure.
25 The residue formed was chromatographed on silica gel with a 95:5:1 v/v/v mixture of dichloromethane, methanol and saturated aqueous ammonia as the eluent giving 37.3 mg (74.5%) Methyl-(2-methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine as light brown solid, MS: 458 (MH^+).

In analogy to the method described in example 52.1, primary or secondary amines were
30 reacted with halo-heterocyclic compounds in N,N-dimethylformamide to yield products as listed in the following table. The compounds were obtained as colorless or light brown oils or solids.

Example	Product	MS MH ⁺	Primary or Secondary Amine	Halo-Heterocycle
52.2	Methyl-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-pyrimidin-4-yl-amine	444	Methyl-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine	4-Chloro-pyrimidine [PCT Int. Appl. (1998), WO9821188A1]
52.3	(2-Methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine	444	2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethylamine	4-Chloro-2-methyl-pyrimidine
52.4	{2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-pyrimidin-4-yl-amine	430	2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethylamine	4-Chloro-pyrimidine
52.5	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-(2-methyl-pyrimidin-4-yl)-amine	496, 1Br	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine	4-Chloro-2-methyl-pyrimidine
52.6	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-pyrimidin-4-yl-amine	482, 1Br	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine	4-Chloro-pyrimidine
52.7	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-(2-methyl-pyrimidin-4-yl)-amine	482, 1Br	4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butylamine	4-Chloro-2-methyl-pyrimidine

52.8	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-pyrimidin-4-yl-amine	468, 1Br	4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butylamine	4-Chloro-pyrimidine
52.9	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-pyridin-4-yl-amine	481 1Br	{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine	4-Chloro-pyridine

Example 53

53.1

- A solution of 100 mg (0.220 mmol) 6-(4-Bromo-butoxy)-3-(4-bromo-phenyl)-2-methyl-benzo[b]thiophene (example 27.5), 0.024 g (0.35 mmol) of imidazole and 0.0144 g (0.330 mmol) sodium hydride (55% suspension in oil) in 0.5 ml of N,N-dimethylformamide was stirred for 1 hour at room temperature. The reaction mixture was then poured into 30 ml of ice-water and extracted 3 times with 10 ml of ether. The combined ether phases were washed with brine, dried over magnesium sulfate and evaporated under reduced pressure.
- The residue formed was chromatographed on silica gel with a 9:1 v/v mixture of dichloromethane and methanol as the eluent yielding 84.2 mg (86.7%) 1-{4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl}-1H-imidazole as colorless oil, MS: 441 (MH⁺, 1Br).

53.2

- In analogy to the procedure described in example 53.1, Methanesulfonic acid 2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl ester (example 49.2) was reacted with imidazole and sodium hydride in N,N-dimethylformamide at room temperature to yield 1-{2-[2-Methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-1H-imidazole as colorless oil, MS: 403 (MH⁺).

Example A

Tablets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per tablet</u>
Compound of formula I	10.0 - 100.0 mg
Lactose	125.0 mg
Maize starch	75.0 mg
Talc	4.0 mg
Magnesium stearate	1.0 mg

5

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula I	25.0 mg
Lactose	150.0 mg
Maize starch	20.0 mg
Talc	5.0 mg


Example C

10 Injection solutions can have the following composition:

Compound of formula I	3.0 mg
Gelatine	150.0 mg
Phenol	4.7 mg
Water for injection solutions	ad 1.0 ml

Claims

1. Compounds of formula (I)




wherein

- 5 U is O or a lone pair,
- Y N, or CR¹,
- Z is S, or S(O₂),
- V is a) -CH=CH-, and m and n are 0, or
 b) -CH₂-, and m+n ≤ 2, or
 10 c) O or NR², and m is 1 to 6, n is 1 to 6, m+n ≤ 7, or m is 1 to 3, n is 0, or
 d) S, and m is 1 to 7, n is 0 to 6, m+n ≤ 7, or
 e) -C≡C-, and m is 0 to 7, n is 0 to 7, m+n is ≤ 7,
- Q is cycloalkyl, cycloalkyl-lower-alkyl, phenyl optionally substituted by 1 to 3
 substituents independently selected from the group as defined for R³, or an
 15 alkyl, alkenyl or alkadienyl group optionally substituted by OH,
- A¹ is hydrogen, lower-alkenyl, or lower-alkyl optionally substituted with hydroxy,
 lower-alkoxy, or thio-lower-alkoxy,
- A² is cycloalkyl, cycloalkyl-lower-alkyl, lower-alkenyl, lower-alkinyl, heteroaryl, or
 lower-alkyl optionally substituted with halogen, hydroxy, lower-alkoxy, or
 20 thio-lower-alkoxy, or
- A¹ and A² are bonded to each other to form a ring and -A¹-A²- is lower-alkylene, or
 lower-alkenylene, optionally substituted by R⁴, in which one -CH₂- group of
 -A¹-A²- can optionally be replaced by a NR⁵, S, or O, or -A¹-A²- is
 -CH=N=CH=CH- which can optionally be substituted by lower-alkyl,
- 25 A³ and A⁴ independently from each other are hydrogen or lower-alkyl, or

- A^3 and A^4 are bonded to each other to form a ring together with the carbon atom to which they are attached and $-A^3-A^4-$ is $-(CH_2)_{2-5}-$ which can optionally be mono- or multiply-substituted by lower-alkyl,
- X is hydrogen or one or more optional halogen and/or lower-alkyl substituents,
- 5 R^1 is hydrogen or lower-alkyl,
- R^2 is hydrogen or lower-alkyl,
- R^3 is halogen, $N(R^6, R^7)$, piperidyl, piperazinyl, piperazinyl-ethanone, morpholinyl, $CONH_2$, CN, NO_2 , CF_3 , OH, lower-alkoxy, thio-lower-alkoxy, or is lower-alkyl, lower-alkenyl, or lower-alkinyl, optionally substituted with OH, SH or $N(R^8, R^9)$,
- 10 R^4 is hydroxy, lower-alkyl, lower-alkoxy, or thio-lower-alkoxy,
- R^5 is hydrogen, lower-alkyl, or lower-alkyl-carbonyl,
- R^6 and R^7 independently from each other are selected from the group consisting of hydrogen, lower-alkyl, phenyl and benzyl,
- 15 R^8 and R^9 independently from each other are selected from the group consisting of hydrogen and lower-alkyl,

and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

2. Compounds of formula (I) according to claim 1



20 wherein

- U is O or a lone pair,
- Y is N, or CR^1 ,
- Z is S, or $S(O_2)$,
- V is a) $-CH=CH-$, and m and n are 0, or
- 25 b) $-CH_2-$, and $m+n \leq 2$, or
- c) O or NR^2 , and m is 1 to 6, n is 1 to 6, $m+n \leq 7$, or m is 1 to 3, n is 0, or
- d) S, and m is 1 to 7, n is 0 to 6, $m+n \leq 7$, or

- e) $\text{-C}\equiv\text{C-}$, and m is 0 to 7, n is 0 to 7, $m+n$ is ≤ 7 ,
- Q is cycloalkyl, cycloalkyl-lower-alkyl, phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R^3 , or an alkyl, alkenyl or alkadienyl group optionally substituted by OH,
- 5 A^1 is hydrogen, lower-alkenyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy,
- A^2 is cycloalkyl, cycloalkyl-lower-alkyl, lower-alkenyl, lower-alkinyl, or lower-alkyl optionally substituted with hydroxy, lower-alkoxy, or thio-lower-alkoxy, or
- 10 A^1 and A^2 are bonded to each other to form a ring and $\text{-A}^1\text{-A}^2\text{-}$ is lower-alkylene, or lower-alkenylene, optionally substituted by R^4 , in which one $\text{-CH}_2\text{-}$ group of $\text{-A}^1\text{-A}^2\text{-}$ can optionally be replaced by a NR^5 , S, or O,
- A^3 and A^4 independently from each other are hydrogen or lower-alkyl,
- X is hydrogen or one or more optional halogen and/or lower-alkyl substituents,
- 15 R^1 is hydrogen or lower-alkyl,
- R^2 is hydrogen or lower-alkyl,
- R^3 is halogen, NH_2 , $\text{N}(\text{lower-alkyl})_2$, CONH_2 , CN, NO_2 , CF_3 , OH, lower-alkoxy, thio-lower-alkoxy, or is lower-alkyl, lower-alkenyl, or lower-alkinyl, optionally substituted with OH, SH or NH_2 ,
- 20 R^4 is hydroxy, lower-alkyl, lower-alkoxy, or thio-lower-alkoxy,
- R^5 is hydrogen, lower-alkyl, or lower-alkyl-carbonyl,

and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

3. Compounds according to any of claims 1 to 2, wherein U is a lone pair.
4. Compounds according to any of claims 1 to 3, wherein V is O, m is 1 to 6,
- 25 n is 1 to 6, $m+n \leq 7$.
5. Compounds according to any of claims 1 to 3, wherein V is O, m is 1 to 3, n is 0.
6. Compounds according to any of claims 1 to 3, wherein V is $\text{-C}\equiv\text{C-}$, m is 0 to 2, n is 0.

7. Compounds according to any of claims 1 to 3, wherein V is $-\text{CH}_2-$ and $m+n \leq 2$.
8. Compounds according to any of claims 1 to 7, wherein A^1 is hydrogen, methyl, or ethyl optionally substituted with hydroxy or methoxy.
- 5 9. Compounds according to any of claims 1 to 8, wherein A^2 is methyl, n-propyl, i-propyl, 2-propenyl, 2-propinyl, 1,1-dimethyl-2-propinyl, cyclopropyl-methylene, or ethyl optionally substituted with hydroxy or methoxy.
- 10 10. Compounds according to any of claims 1 to 9, wherein A^2 is methyl, ethyl, 2-hydroxy-ethyl, 2-methoxy-ethyl, n-propyl, 2-propenyl, or cyclopropyl-methylene.
- 11 11. Compounds according to any of claims 1 to 8, wherein A^2 is 2-Methyl-pyrimidin-4-yl.
- 12 12. Compounds according to any of claims 1 to 7, wherein A^1 and A^2 are bonded to each other to form a ring and $-\text{A}^1-\text{A}^2-$ is lower-alkylene, optionally substituted by R^4 , in which one $-\text{CH}_2-$ group of $-\text{A}^1-\text{A}^2-$ can optionally be replaced by NR^5 or O, wherein R^4 is
15 hydroxy, and R^5 is lower-alkyl or lower-alkyl-carbonyl, or $-\text{A}^1-\text{A}^2-$ is $-\text{CH}=\text{N}=\text{CH}=\text{CH}-$.
13. Compounds according to claim 12, wherein R^5 is methyl.
14. Compounds according to claim 12, wherein $-\text{A}^1-\text{A}^2-$ is $-(\text{CH}_2)_5-$.
15. Compounds according to any of claims 1 to 14, wherein A^3 is hydrogen.
16. Compounds according to any of claims 1 to 15, wherein A^4 is hydrogen.
- 20 17. Compounds according to any of claims 1 to 14, wherein A^3 is methyl.
18. Compounds according to any of claims 1 to 15, wherein A^4 is methyl.
19. Compounds according to any of claims 1 to 18, wherein Y is CR^1 and R^1 is methyl.
20. Compounds according to any of claims 1 to 19, wherein Y is N.
- 25 21. Compounds according to any of claims 1 to 20, wherein Z is S.
22. Compounds according to any of claims 1 to 20, wherein Z is SO_2 .

23. Compounds according to any of claims 1 to 22, wherein Q is phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R³, or an alkyl or alkenyl group optionally substituted by OH, wherein R³ is as defined in claim 1.
- 5 24. Compounds according to any of claims 1 to 23, wherein Q is phenyl optionally substituted by 1 to 3 substituents independently selected from the group as defined for R³, wherein R³ is fluorine, chlorine, bromine, or CF₃.
25. Compounds according to any of claims 1 to 24, wherein Q is 4-chloro-phenyl, 4-bromo-phenyl, or 4-trifluoromethyl-phenyl.
- 10 26. Compounds according to any of claims 1 to 25, wherein X is hydrogen.
27. A compound according to any of claims 1 to 26, selected from the group consisting of
- 2-({5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol,
 2-(Ethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amino)-
 15 ethanol,
 2-[[4-[3-(4-Bromo-phenyl)-2-methyl-benzo[b]thiophen-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol,
 {5-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-dimethyl-amine,
 {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-
 20 ethyl-(2-methoxy-ethyl)-amine,
 3-(4-Bromo-phenyl)-6-[4-(4-methyl-piperazin-1-yl)-butoxy]-benzo[d]isothiazole,
 {4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-diethyl-amine, and
 Allyl-methyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-
 25 amine,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.
28. A compound according to any of claims 1 to 26, selected from the group consisting of
- 2-[[3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-propyl]-ethyl-amino]-ethanol,
 30 Dimethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amine,
 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-ethyl-amino]-ethanol,
 6-(3-Azetidin-1-yl-prop-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole,
 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl}-amino)-

- ethanol,
 2-[[4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol,
 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-
 5 cyclopropylmethyl-methyl-amine, and
 Dimethyl-{5-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-amine,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

29. A compound according to any of claims 1 to 26, selected from the group consisting of
 10 {5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-diethyl-amine,
 2-({5-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yl]-pent-4-ynyl}-ethyl-amino)-ethanol,
 Allyl-{4-[3-(4-bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl}-methyl-amine,
 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-methyl-
 15 propyl-amine,
 2-[[4-[3-(4-Bromo-phenyl)-2-methyl-1,1-dioxo-1H-11 6-benzo[b]thiophen-6-yloxy]-butyl]-(2-hydroxy-ethyl)-amino]-ethanol,
 6-(5-Azetidin-1-yl-pent-1-ynyl)-3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazole, and
 2-(Ethyl-{4-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-but-3-ynyl}-amino)-
 20 ethanol,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.


30. A compound according to any of claims 1 to 26, selected from the group consisting of
 {4-[3-(4-Chloro-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl}-ethyl-(2-
 25 methoxy-ethyl)-amine,
 (4-[3-(4-Bromo-phenyl)-1,1-dioxo-1H-11 6-benzo[d]isothiazol-6-yloxy]-butyl)-dimethyl-amine, and
 2-((4-[3-(4-Chloro-phenyl)-benzo[d]isothiazol-6-yloxy]-butyl)-methyl-amino)-ethanol,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.

- 30 31. A compound according to any of claims 1 to 26, selected from the group consisting of
 3-(4-Bromo-phenyl)-6-(3-piperidin-1-yl-propoxy)-benzo[d]isothiazole,
 2-(Ethyl-{4-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-butyl}-amino)-ethanol,

- 2-(Ethyl-{1-methyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl}-amino)-ethanol,
 Ethyl-(2-methoxy-ethyl)-{4-[2-methyl-1,1-dioxo-3-(4-trifluoromethyl-phenyl)-1H-1l 6-benzo[b]thiophen-6-yloxy]-butyl}-amine,
 5 {4-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-butyl}-ethyl-(2-methoxy-ethyl)-amine,
 2-(Ethyl-{3-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yl]-propyl}-amino)-ethanol,
 2-((1,1-Dimethyl-3-[3-(4-trifluoromethyl-phenyl)-benzo[d]isothiazol-6-yl]-prop-2-ynyl)-ethyl-amino)-ethanol,
 10 2-((3-[3-(4-Bromo-phenyl)-benzo[d]isothiazol-6-yl]-1,1-dimethyl-prop-2-ynyl)-ethyl-amino)-ethanol,
 Methyl-(2-methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-benzo[b]thiophen-6-yloxy]-ethyl}-amine, and
 (2-Methyl-pyrimidin-4-yl)-{2-[2-methyl-3-(4-trifluoromethyl-phenyl)-
 15 benzo[b]thiophen-6-yloxy]-ethyl}-amine,
 and pharmaceutically acceptable salts and/or pharmaceutically acceptable esters thereof.


32. A process for the manufacture of compounds according to any of claims 1 to 31, which process comprises

a) reacting a compound of formula (II)




with a compound $(A^1, A^2, U)N-C(A^3, A^4)-(CH_2)_m-M$, wherein V is O, S or NR^2 , M is mesylate, tosylate, triflate, Cl, Br or I, and X, Y, Z, Q, U, A^1 , A^2 , A^3 , A^4 , m, n and R^2 have the significances given in claim 1, or wherein HV is mesylate, tosylate, triflate, Cl, Br or I, and M is OH, SH or NHR^2 , and R^2 has the significance given in claim 1,

25 or b) reacting a compound of formula (III)




with a compound NHA^1A^2 , wherein M is mesylate, tosylate, triflate, Cl, Br or I, and $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4, \text{V}, \text{X}, \text{Y}, \text{Z}, \text{Q}, \text{m}$ and n are as defined in claim 1,

or c) reacting a compound of formula (IV)




with a compound $(\text{A}^1, \text{A}^2, \text{U})\text{N}-\text{C}(\text{A}^3, \text{A}^4)-(\text{CH}_2)_m-\text{C}\equiv\text{CH}$, wherein X, Y, Z, Q, U, $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4$ and m are as defined in claim 1,

or d) reacting a compound of formula (V)



with a compound $(\text{A}^1, \text{A}^2, \text{U})\text{N}-\text{C}(\text{A}^3, \text{A}^4)-(\text{CH}_2)_m-\text{M}$, wherein M is mesylate, tosylate, Cl, Br or I, and $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4, \text{U}, \text{X}, \text{Y}, \text{Z}, \text{Q}, \text{m}$ and n are as defined in claim 1,

or e) hydrogenating a compound of formula (VI)



wherein V is $-\text{C}\equiv\text{C}-$, and $\text{A}^1, \text{A}^2, \text{A}^3, \text{A}^4, \text{U}, \text{X}, \text{Y}, \text{Z}, \text{Q}, \text{m}$ and n are as defined in claim 1,

and optionally converting a compound according to any of claims 1 to 31, wherein U is a lone pair, to a corresponding compound wherein U is O.

33. Compounds according to any of claims 1 to 31 when manufactured by a process according to claim 32.

5 34. Pharmaceutical compositions comprising a compound according to any of claims 1 to 31 and a pharmaceutically acceptable carrier and/or adjuvant.

35. Compounds according to any of claims 1 to 31 for use as therapeutic active substances.

10 36. Compounds according to any of claims 1 to 31 for use as therapeutic active substances for the treatment and/or prophylaxis of diseases which are associated with OSC.

15 37. A method for the treatment and/or prophylaxis of diseases which are associated with OSC such as hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones, tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance and diabetes, which method comprises administering a compound according to any of claims 1 to 31 to a human being or animal.

38. The use of compounds according to any of claims 1 to 31 for the treatment and/or prophylaxis of diseases which are associated with OSC.

20 39. The use of compounds according to any of claims 1 to 31 for the treatment and/or prophylaxis of hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones, tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance, and diabetes.

25 40. The use of compounds according to any of claims 1 to 31 for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are associated with OSC.

30 41. The use of compounds according to any of claims 1 to 31 for the preparation of medicaments for the treatment and/or prophylaxis of hypercholesterolemia, hyperlipemia, arteriosclerosis, vascular diseases, mycoses, parasite infections, gallstones,

tumors and/or hyperproliferative disorders, and/or treatment and/or prophylaxis of impaired glucose tolerance and diabetes.

42. The novel compounds, processes and methods as well as the use of such compounds substantially as described hereinbefore.

5

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 01/12451

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D333/54 C07D275/04 C07D409/12 C07D417/12 A61K31/381
A61P3/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 778 271 A (HOFFMANN LA ROCHE) 11 June 1997 (1997-06-11) page 14, line 30 -page 16, line 55; claims 1,5	1-41
X	US 4 410 539 A (CROSS PETER E ET AL) 18 October 1983 (1983-10-18) claim 1; examples 12,14,46,47,56,57 -/-	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

22 January 2002

Date of mailing of the international search report

29/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seelmann, I

INTERNATIONAL SEARCH REPORT

Intern Application No

PCT/EP 01/12451

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CORSS P E ET AL: "SELECTIVE THROMBOXANE SYNTHETASE INHIBITORS. 3. 1H-IMIDAZOL-1-YL-SUBSTITUTED BENZOUBFURAN-, BENZOUTHIOPHENE-. ANS INDOLE-2- AND -3-CARBOXYLIC ACIDS" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 29, no. 9, 1986, pages 1637-1643, XP002066696 ISSN: 0022-2623 compounds 42 and 76	1
A	US 5 994 371 A (CULLINAN GEORGE J ET AL) 30 November 1999 (1999-11-30) column 4, line 1 - line 57	1-41

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 42

Present claim 42 relates to an extremely large number of possible compounds and methods. In fact, the claim is formulated in such a general way that a lack of clarity within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claim impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely claims 1-41.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT
information on patent family members

International Application No

PCT/EP 01/12451

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0778271	A	11-06-1997	BR 9605906 A	18-08-1998
			CA 2190708 A1	09-06-1997
			CN 1161328 A ,B	08-10-1997
			EP 0778271 A2	11-06-1997
			JP 9176123 A	08-07-1997
			TR 970475 A2	21-06-1997
			US 5856503 A	05-01-1999
US 4410539	A	18-10-1983	DE 3165703 D1	27-09-1984
			DK 466681 A	24-04-1982
			EP 0050957 A1	05-05-1982
			GB 2101115 A	12-01-1983
			GR 75101 A1	13-07-1984
			IE 51650 B1	21-01-1987
			JP 57181084 A	08-11-1982
			JP 63035630 B	15-07-1988
			US 4551468 A	05-11-1985
US 5994371	A	30-11-1999	US 5482949 A	09-01-1996
			US 5605924 A	25-02-1997
			US 5599833 A	04-02-1997
			US 5798351 A	25-08-1998
			AT 180479 T	15-06-1999
			AU 670177 B2	04-07-1996
			AU 5786394 A	22-09-1994
			BR 9401183 A	01-11-1994
			CN 1097420 A	18-01-1995
			CZ 9400578 A3	15-12-1994
			DE 69418639 D1	01-07-1999
			DE 69418639 T2	07-10-1999
			DK 617030 T3	08-11-1999
			EP 0617030 A1	28-09-1994
			ES 2132339 T3	16-08-1999
			FI 941262 A	20-09-1994
			GR 3030317 T3	30-09-1999
			HU 70549 A2	30-10-1995
			JP 6321937 A	22-11-1994
			NO 940940 A	20-09-1994
			NZ 260085 A	21-12-1995
			PH 30177 A	21-01-1997
			SI 617030 T1	31-10-1999
			ZA 9401786 A	14-09-1995